

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫ
Satbayev University

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ
НАУК РЕСПУБЛИКИ
КАЗАХСТАН
Satbayev University

N E W S

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF
KAZAKHSTAN
Satbayev University

SERIES
OF GEOLOGY AND TECHNICAL SCIENCES

2 (452)
MARCH – APRIL 2022

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы «ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы» ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Бас редактор

ЖҰРЫНОВ Мұрат Жұрынұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Қазақстан Республикасы Ұлттық Ғылым академиясының президенті, АҚ «Д.В. Сокольский атындағы отын, катализ және электрохимия институтының» бас директоры (Алматы, Қазақстан) **Н = 4**

АБСАДЫКОВ Бахыт Нарикбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА жауапты хатшысы, А.Б. Бектұров атындағы химия ғылымдары институты (Алматы, Қазақстан) **Н = 5**

Редакциялық алқа:

ӘБСАМЕТОВ Мәліс Құдысұлы (бас редактордың орынбасары), геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, «У.М. Ахмедсафина атындағы гидрогеология және геоэкология институтының» директоры (Алматы, Қазақстан) **Н = 2**

ЖОЛТАЕВ Герой Жолтайұлы (бас редактордың орынбасары), геология-минералогия ғылымдарының докторы, профессор, Қ.И. Сатпаев атындағы геология ғылымдары институтының директоры (Алматы, Қазақстан) **Н = 2**

СНОУ Дэниел, Ph.D, қауымдастырылған профессор, Небраска университетінің Су ғылымдары зертханасының директоры (Небраска штаты, АҚШ) **Н = 32**

ЗЕЛЬТМАН Реймар, Ph.D, табиғи тарих мұражайының Жер туралы ғылымдар бөлімінде петрология және пайдалы қазбалар кен орындары саласындағы зерттеулердің жетекшісі (Лондон, Англия) **Н = 37**

ПАНФИЛОВ Михаил Борисович, техника ғылымдарының докторы, Нанси университетінің профессоры (Нанси, Франция) **Н = 15**

ШЕН Пин, Ph.D, Қытай геологиялық қоғамының тау геологиясы комитеті директорының орынбасары, Американдық экономикалық геологтар қауымдастығының мүшесі (Пекин, Қытай) **Н = 25**

ФИШЕР Аксель, Ph.D, Дрезден техникалық университетінің қауымдастырылған профессоры (Дрезден, Берлин) **Н = 6**

КОНТОРОВИЧ Алексей Эмильевич, геология-минералогия ғылымдарының докторы, профессор, РФА академигі, А.А. Трофимука атындағы мұнай-газ геологиясы және геофизика институты (Новосибирск, Ресей) **Н = 19**

АГАБЕКОВ Владимир Енокович, химия ғылымдарының докторы, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры (Минск, Беларусь) **Н = 13**

КАТАЛИН Стефан, Ph.D, Дрезден техникалық университетінің қауымдастырылған профессоры (Дрезден, Берлин) **Н = 20**

СЕЙТМҰРАТОВА Элеонора Юсуповна, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА корреспондент-мүшесі, Қ.И. Сатпаев атындағы Геология ғылымдары институты зертханасының меңгерушісі (Алматы, Қазақстан) **Н = 11**

САҒЫНТАЕВ Жанай, Ph.D, қауымдастырылған профессор, Назарбаев университеті (Нұр-Сұлтан, Қазақстан) **Н = 11**

ФРАТТИНИ Паоло, Ph.D, Бикокк Милан университеті қауымдастырылған профессоры (Милан, Италия) **Н = 28**

«ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктеуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және қоғамдық даму министрлігінің Ақпарат комитетінде 29.07.2020 ж. берілген № **KZ39VPY00025420** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *геология, мұнай және газды өңдеудің химиялық технологиялары, мұнай химиясы, металдарды алу және олардың қосындыларының технологиясы.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекен-жайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., тел.: 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2022

Типографияның мекен-жайы: «Аруна» ЖК, Алматы қ., Мұратбаев көш., 75.

Главный редактор

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик НАН РК, президент Национальной академии наук Республики Казахстан, генеральный директор АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского» (Алматы, Казахстан) **Н = 4**

АБСАДЫКОВ Бахыт Нарикбаевич, доктор технических наук, профессор, ответственный секретарь НАН РК, Институт химических наук им. А.Б. Бектурова (Алматы, Казахстан) **Н = 5**

Редакционная коллегия:

АБСАМЕТОВ Малис Кудысович, (заместитель главного редактора), доктор геологоминералогических наук, профессор, академик НАН РК, директор Института гидрогеологии и геоэкологии им. У.М. Ахмедсафина (Алматы, Казахстан) **Н = 2**

ЖОЛТАЕВ Герой Жолтаевич, (заместитель главного редактора), доктор геологоминералогических наук, профессор, директор Института геологических наук им. К.И. Сатпаева (Алматы, Казахстан) **Н=2**

СНОУ Дэниел, Ph.D, ассоциированный профессор, директор Лаборатории водных наук университета Небраска, США) **Н = 32**

ЗЕЛЬТМАН Реймар, Ph.D, руководитель исследований в области петрологии и месторождений полезных ископаемых в Отделе наук о Земле Музея естественной истории (Лондон, Англия) **Н = 37**

ПАНФИЛОВ Михаил Борисович, доктор технических наук, профессор Университета Нанси (Нанси, Франция) **Н=15**

ШЕН Пин, Ph.D, заместитель директора Комитета по горной геологии Китайского геологического общества, член Американской ассоциации экономических геологов (Пекин, Китай) **Н = 25**

ФИШЕР Аксель, ассоциированный профессор, Ph.D, технический университет Дрезден (Дрезден, Берлин) **Н = 6**

КОНТОРОВИЧ Алексей Эмильевич, доктор геолого-минералогических наук, профессор, академик РАН, Институт нефтегазовой геологии и геофизики им. А.А. Трофимука СО РАН (Новосибирск, Россия) **Н = 19**

АГАБЕКОВ Владимир Енокович, доктор химических наук, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь) **Н = 13**

КАТАЛИН Стефан, Ph.D, ассоциированный профессор, Технический университет (Дрезден, Берлин) **Н = 20**

СЕЙТМУРАТОВА Элеонора Юсуповна, доктор геолого-минералогических наук, профессор, член-корреспондент НАН РК, заведующая лабораторией Института геологических наук им. К.И. Сатпаева (Алматы, Казахстан) **Н=11**

САГИНТАЕВ Жанай, Ph.D, ассоциированный профессор, Назарбаев университет (Нурсултан, Казахстан) **Н = 11**

ФРАТТИНИ Паоло, Ph.D, ассоциированный профессор, Миланский университет Бикокк (Милан, Италия) **Н = 28**

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № **KZ39VPY00025420**, выданное 29.07.2020 г.

Тематическая направленность: *геология, химические технологии переработки нефти и газа, нефтехимия, технологии извлечения металлов и их соединений.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, оф. 219, тел.: 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2022

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editorial chief

ZHURINOV Murat Zhurinovich, doctor of chemistry, professor, academician of NAS RK, president of the National Academy of Sciences of the Republic of Kazakhstan, general director of JSC “Institute of fuel, catalysis and electrochemistry named after D.V. Sokolsky» (Almaty, Kazakhstan) **H = 4**

ABSADYKOV Bakhyt Narikbaevich, doctor of technical sciences, professor, executive secretary of NAS RK, Bekturov Institute of chemical sciences (Almaty, Kazakhstan) **H = 5**

Editorial board:

ABSAMETOV Malis Kudysovich, (deputy editor-in-chief), doctor of geological and mineralogical sciences, professor, academician of NAS RK, director of the Akhmedsafin Institute of hydrogeology and hydrophysics (Almaty, Kazakhstan) **H=2**

ZHOLTAEV Geroy Zholtaevich, (deputy editor-in-chief), doctor of geological and mineralogical sciences, professor, director of the institute of geological sciences named after K.I. Satpayev (Almaty, Kazakhstan) **H=2**

SNOW Daniel, Ph.D, associate professor, director of the laboratory of water sciences, Nebraska University (Nebraska, USA) **H = 32**

ZELTMAN Reymer, Ph.D, head of research department in petrology and mineral deposits in the Earth sciences section of the museum of natural history (London, England) **H = 37**

PANFILOV Mikhail Borisovich, doctor of technical sciences, professor at the Nancy University (Nancy, France) **H=15**

SHEN Ping, Ph.D, deputy director of the Committee for Mining geology of the China geological Society, Fellow of the American association of economic geologists (Beijing, China) **H = 25**

FISCHER Axel, Ph.D, associate professor, Dresden University of technology (Dresden, Germany) **H=6**

KONTOROVICH Aleksey Emilievich, doctor of geological and mineralogical sciences, professor, academician of RAS, Trofimuk Institute of petroleum geology and geophysics SB RAS (Novosibirsk, Russia) **H = 19**

AGABEKOV Vladimir Enokovich, doctor of chemistry, academician of NAS of Belarus, honorary director of the Institute of chemistry of new materials (Minsk, Belarus) **H = 13**

KATALIN Stephan, Ph.D, associate professor, Technical university (Dresden, Berlin) **H = 20**

SEITMURATOVA Eleonora Yusupovna, doctor of geological and mineralogical sciences, professor, corresponding member of NAS RK, head of the laboratory of the Institute of geological sciences named after K.I. Satpayev (Almaty, Kazakhstan) **H=11**

SAGINTAYEV Zhanay, Ph.D, associate professor, Nazarbayev University (Nursultan, Kazakhstan) **H = 11**

FRATTINI Paolo, Ph.D, associate professor, university of Milano-Bicocca (Milan, Italy) **H = 28**

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA «National Academy of Sciences of the Republic of Kazakhstan» (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan **No. KZ39VPY00025420**, issued 29.07.2020.

Thematic scope: *geology, chemical technologies for oil and gas processing, petrochemistry, technologies for extracting metals and their connections.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, Almaty, 050010, tel. 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2022

Address of printing house: ST «Aruna», 75, Muratbayev str, Almaty.

NEWS of the National Academy of Sciences of the Republic of Kazakhstan
SERIES OF GEOLOGY AND TECHNICAL SCIENCES
ISSN 2224-5278

Volume 2, Number 452 (2022), 33-48

<https://doi.org/10.32014/2022.2518-170X.158>

UDC 550.3(574)

IRSTI 38.57

Zh.K. Aidarbekov*, S.A. Istekova

The Satbayev University, Almaty, Kazakhstan.

E-mail: zh.aidarbekov@gmail.com

**CLASSIFICATION OF GEOPHYSICAL FIELDS IN THE STUDY
OF GEOLOGICAL AND STRUCTURAL FEATURES OF THE
ZHEZKAZGAN ORE DISTRICT**

Abstract. As a result of generalization and analysis of complex geological and geophysical data, the geophysical studies were shown in application to structural mapping of ore-prospective areas and detailed study of ore-controlling complexes of cupreous sandstones in the Zhezkazgan ore district of Central Kazakhstan. Basing on modern means of registration, processing and interpretation of magnetic, gravity and seismic exploration, the ore geophysics proved its efficacy in studies of tectonic structure of ore areas, localisation and identification of the morphology of differentiated intrusive massifs at depth, detection and deep mapping of tectonic faults, defining the ore-controlling structures of sedimentary and effusive-sedimentary folded complexes as well as in determination of the spatial position of ore-bearing zones. The deep structure of the Zhezkazgan syncline on sides of the Sary-Oba rise differs sharply, fixing the simple structure of the eastern part and the complex folded one of the western part. The deposits of the Zhezkazgan and overlying zhidelisai and kingir suites represent a conforming continuous section, in places, there are local disconformities over faults with repeated tectonic displacements, lying on sediments of the taskuduk suite. Taking into account the rather complex geological structure of the given ore district and the significant differentiation of the composing rocks, ores and near-ore hosting strata, further identification of additional criteria for localization of copper mineralization is possible with the integration of detailed geological and geophysical methods, geochemical and physical-chemical studies of rocks and ores.

Key words: cupreous sandstones, geophysical fields, interpretation, ore-bearing zones.

Айдарбеков Ж.К.*, Истекова С.А.

Сәтбаев атындағы университет, Алматы, Қазақстан.

E-mail: zh.aidarbekov@gmail.com

ЖЕЗҚАЗҒАН КЕНДІ АЙМАҒЫНЫҢ ГЕОЛОГИЯЛЫҚ ҚҰРЫЛЫС ЕРЕКШЕЛІКТЕРІН ЗЕРТТЕУ БАРЫСЫНДАҒЫ ГЕОФИЗИКАЛЫҚ ӨРІСТЕРІН КЛАССИФИКАЦИЯЛАУ

Аннотация. Кешенді геологиялық-геофизикалық мәліметтерді қорыту және талдау нәтижесінде Орталық Қазақстанның Жезқазған кен аймағындағы кен перспективалы жабық аумақтардың құрылымдық картасын жасау және мыс құмтастарының рудалық бақылау кешендерін егжей-тегжейлі зерделеу үшін геофизикалық зерттеулер жүргізу мүмкіндігі пайда болды. Магниттік, гравитациялық, сейсмикалық барлауды тіркеудің, өңдеудің және интерпретациялаудың заманауи құралдарын пайдалану негізінде кенді аймақтардың құрылымдық-тектоникалық құрылымын зерттеуде кен геофизикасының тиімділігі көрсетілді; тереңдікте дифференциалды интрузивті массивтердің морфологиясын локализациялау және анықтау; тектоникалық бұзылуларды анықтау және терең картаға түсіру, шөгінді және эффузивті-шөгінді қатпарлы кешендердегі кенді реттейтін құрылымдарды анықтау; кенді аймақтардың кеңістіктегі орнын анықтау. Жезқазған синклиналының терең құрылымы Сарыоба жыртылысының әр бетінде күрт ерекшеленеді. Зерттелетін кенді аймақтың айтарлықтай дифференциациясын ескере отырып, мыс минерализациясының одан әрі анықтау егжей-тегжейлі геологиялық, геофизикалық әдістерді геохимиялық және физика-химиялық зерттеулермен біріктіру арқылы мүмкін болады.

Түйін сөздер: мыс құмтастар, геофизикалық өрістер, интерпретация, кенді аймақтар.

Ж.К. Айдарбеков*, С.А.Истекова

Satbayev University, Алматы, Қазақстан.

E-mail: zh.aidarbekov@gmail.com

РАЙОНИРОВАНИЕ ГЕОФИЗИЧЕСКИХ ПОЛЕЙ ПРИ ИЗУЧЕНИИ ГЕОЛОГО-СТРУКТУРНЫХ ОСОБЕННОСТЕЙ ЖЕЗКАЗГАНСКОГО РУДНОГО РАЙОНА

Аннотация. В результате обобщения и анализа комплексных геолого-геофизических данных показана возможность геофизических исследова-

ний для структурного картирования рудоперспективных закрытых площадей и детального изучения рудоконтролирующих комплексов медистых песчаников в Жезказганском рудном районе Центрального Казахстана. На основе применения современных средств регистрации, обработки и интерпретации магнито-, грави- и сейсморазведки показана эффективность рудной геофизики при изучении структурно-тектонического строения рудных районов; локализации и выявлении морфологии дифференцированных интрузивных массивов на глубине; обнаружения и глубинного картирования тектонических разломов, выделения рудоконтролирующих структур в осадочных и эффузивно-осадочных складчатых комплексах; определении пространственного положения рудоносных зон. Глубинное строение Жезказганской синклинали резко различается по разные стороны Сарыобинского взброса, при этом фиксируется простое строение восточной части и сложное складчатое для западной половины. Отложения жезказганской и вышележащих жиделисайской и кингирской свит представляют собой согласный непрерывный разрез, местами отмечаются местные несогласия над разломами с неоднократно повторявшимися тектоническими смещениями, залегающий на отложениях таскудукской свиты. Учитывая достаточно сложное геологическое строение исследуемого рудного района и значительную дифференциацию слагающих их пород, руд и околорудных вмещающих толщ, дальнейшее выявление дополнительных критериев локализации медного оруденения представляется возможным при комплексировании детальных геологических и геофизических методов, геохимических и физико-химическим исследований пород и руд.

Ключевые слова: медистые песчаники, геофизические поля, интерпретация, рудоносные зоны.

Introduction. Deposits of cupreous sandstones are one of the leading types in terms of reserves and copper mining in Kazakhstan. The Zhezkazgan ore district includes the unique Zhezkazgan deposit and a number of small copper deposits of the Zhilandy group (It-Aauz, Sary-Oba, Kipshakpai, Karash-Oshak and Zhartas), ore occurrences of the Northern and Southern Akchii, Spassky, Taskuduk, etc. The copper mineralization is associated with deposits of the Taskuduk and Zhezkazgan suites of the Middle and Upper Carboniferous and is confined to layers of their gray-colored sandstones. By formation conditions, the copper mineralization belongs to the stratiform type (Baibatsha A. et.al, 2017).

Until recently, direct search for Zhezkazgan type of ores were confidently carried out only by methods of litho-chemical surveys and the method of induced polarization (IP), mainly, in open areas. Other geophysical methods

of prospecting, including electrical resistance, gravity, magnetic and seismic methods of exploration did not solve this problem, but in search for favourable structures of locked areas they could assist to targeted exploratory drilling (Istekova S.A. 2006) (Uzhkenov B.S. 2011).

In general, the Zhezkazgan ore region was well studied by detailed geological surveys, prospecting and geophysical studies, including a wide range of methods. In different years, magnetic exploration, gravity exploration, electrical exploration, and seismic exploration were implemented in separate sections of the area. The surveys defined the area as a promising one for copper mineralization. For its search at depth, few deep boreholes were drilled (the deepest was 1587.7 meters down), however none of the them have managed to cross the ore-bearing horizons of Zhezkazgan and Tastykuduk suites, while part of the wells have not been tested (Uzhkenov B.S. 2011).

Contemporary means of physical field registration, processing and interpretation allowed to produce high-quality materials and increase efficiency of geophysical methods when solving geological problems at ore deposits.

The high-precision geophysics was applied for both structural mapping of ore-promising areas and detailed study of deep-lying ore-bearing complexes. Generalization and comprehensive analysis of accumulated geophysical information on the Zhezkazgan ore region, with help of modern processing and interpretive geo-information systems, allowed to study in detail the deep-lying ore-bearing complexes and identify additional deep search criteria for stratiform copper mineralization. The territory of the southern immersion of the Western Sary-Oba deposit, covered by loose and Upper Paleozoic sediments, is currently unexplored and remains one of the most promising for replenishing the mineral resource base of the Zhezkazgan region.

Materials and basic methods. The studies were carried out in order to develop search criteria for copper mineralization and to prepare recommendations for exploratory drilling based on geophysical data at the sites of the southern immersion of the Western Sary-Oba deposit within the Zhezkazgan ore district (figure 1).




Figure 1. Layout of deposits and ore occurrences of the Zhezkazgan district (Uzhva V.I. 2011).

Regionally, the study area is located within the Zhezkazgan ore district and covers fragments of the Sarysu-Teniz, Zhezkazgan structural-fomation zone (SFZ), and small volumes of the Konsky and Shagyrlы SFZ. The study area is composed of carbonate-terrigenous deposits of the Carboniferous and Permian systems; in the west part of the district there are terrigenous formations of the Ordovician and Devonian systems along the Eskuly block framing (Zientek M.L. et.al, 2014).

Legend to the layout of the tectonic structure of the Zhilandy ore district:

- boundaries of main structures
- anticipated faults
- 1. deposits
- 2. ore deposits
- reliable faults (berg beames towards the block)
- faults as per geophysical data
- areas of 2012 search and appraisal works

Most of the Zhezkazgan ore district is occupied by the Zhezkazgan syncline, which is a symmetrical structure with the axis dipping to the south in the

meridional direction at an angle of 3-5°. Its western side is steep (at an angle of 70-90°), from the Kokdombak occurrence and to the north has reverse dipping (to the west). The eastern side is flat, dipping at 10-15° to the west. Along the northern side of the syncline, the steepest angles are located to the south of the Eastern Sary-Oba deposit (up to 70-80°); the rock flattens down to 20-12° in the west of the deposit and to 45-30° in its east.

In the northern part of the Zhezkazgan syncline, there is the Northern (Zhilandy) group of copper ore objects, including the deposits of It-Auz, Western and Eastern Sary-Oba, Kipshakpai, Karashoshak and the ore occurrences of Airambai, Pektas, Kokdombak, and oth.

The area of the southern immersion of the Western Sary-Oba deposit within the Zhezkazgan syncline has a meridional strike between the Zhilandy and Zhezkazgan ore fields; is almost by 80% covered by loose formations with a total thickness of up to 30-120m. The indigenous outcrops of Paleozoic rocks are concentrated mainly in the northeastern part of the site and in a small amount are present in the central, southwestern and southeastern parts of the site, where the area of local indigenous outcrops of Late Paleozoic formations does not exceed 1 sq.km. Paleozoic formations are presented by the Zhezkazgan suite of the Middle Upper Carboniferous period, the Zhidelisai suite of the Lower Permian and the Kingir suite of the Lower-Upper Permian period (Cossete P.M et.al, 2014).

The structure of the Zhilandy group of deposits is characterized by complex tectonics, especially within the fields of Western and Eastern Sary-Oba and It-Auz. Sites of the deposits are the zones of maximal stress causing folds and large disjunctive ruptures (Hitzman M.W. et.al, 2010).

Prospects of the planned exploration works in the area of the Zhezkazgan syncline are determined, first of all, by the presence of productive horizons predicted by results of the areal geophysical works and confirmed by exploratory drilling in the past periods. Productive horizons of the Taskuduk suite in the northern part of the Zhezkazgan syncline were revealed during the search and evaluation work carried out by the Zhezkazgan GEE in 1981-1986 (Khailovsky V.N. et al, 1987). The authors established that due to steep rock dipping at the syncline wings, the productive horizons depths of the Taskuduk and Beleuty suites grow sharply down to 1800-1900 m.

In the southeastern part of the Zhezkazgan syncline, in the area of its junction with the Kingir and Uitas brachyanticlines, there are a small deposit Zhartas and an ore occurrence Sorkuduk, both of which are close to the Zhezkazgan type, but are much smaller in scale and with a narrower stratigraphic interval of mineralization (Lyubetskaya L.D. et.al, 2009).

The seismic surveys of 1975-1977 and 1980-1981 performed by the Sarysu and West Teniz seismic parties by the method of reflected waves and CDP

indicated to the spread of productive horizons within the Zhezkazgan syncline. The works resulted in delivery of seismogeological sections and a map of the isodepths for the sole of the productive strata in the central and western parts of the Zhezkazgan syncline.

Research methodology. Additional search criteria for localization of a copper mineralization within the study area were identified using a complex of geophysical methods, including gravity and magnetic exploration, performed in the last decade by geophysical companies applying modern hardware and methodological complexes. The interpretation results obtained by geophysical methods of potential fields were linked to the 2D areal seismic survey data carried out in the area in different periods of time. The main methodological principle was to identify additional search criteria for the localization of copper deposits, from the standpoint of the deep structure: structural-tectonic, stratigraphic and lithological heterogeneity of geological complexes with extensive use of geophysical data (Lyubetskaya L.D. et.al, 2009; Togizov K.S. et.al, 2019).

The following basic localization conditions of Zhezkazgan type copper mineralization contribute to the successful application of structural geophysical methods:

- stratigraphic and lithological control of mineralization, i.e., occurrence of copper mineralization to sandy rocks of the Carboniferous system, namely, to gray medium-coarse-grained sandstones;

- attribution of deposits to peripheral areas of large structures of the first order, having planar dimensions of about 20-30 km; relatively higher copper enrichment of sites, complicated by local folds and ruptures and located along the lines of extended, repeatedly renewed tectonic seams (Balk P.I. et.al, 2016).

Gravity exploration is the main geophysical method applied to studies of deep ore-controlling structure geology. The main parameter affecting the intensity of the gravitational field is the density of rocks which clearly varies in dependence of composition and substances of the geological complexes (Balk P.I et.al, 2017; Cossette P.M. et.al, 2014).

In studies of deep structures, gravimetric maps of observed field of various scales serve as a basis since they contain the most complete information about the whole region and individual target objects. The Zhezkazgan ore region was intensely studied by gravimetric methods. Therefore, deep structures at the target Middle and Lower Paleozoic horizons are provided with sufficient gravimetric data. The territory is fully covered by the 1:200,000 conditional gravimetric survey, and ore areas are covered by surveys scaled 1: 50,000 and larger. The constructed gravimetric maps are of high quality. Maps of the observed field are accompanied by gravitational field transformation maps: regional and local components (filtering by high-frequency bandpass, narrow bandpass and

medium-width filtering), and vertical and horizontal derivative of the Bugey anomaly. Iso-anomal cross-sections were made by 1 and 0.5 mGal.

The gravitational field of the research area is quite differentiated (figure 2).

In the map of the gravitational field, conjunction parts are highlighted for zones with the field's different levels and character. In the western part, the gravitational field is characterized by a submeridional strike of isoanomalos V_{zz} , corresponding to the position of the of the Ulytau-Arganaty structural-formation zone structures. In the southeastern part, the elongated positive and negative anomalies of the northeastern direction are distinguished, spatially coinciding with the Sarysu-Teniz riftogenic structure and the eastern subzone of the Konsky SFZ.

In the central part of the district, there is a band of extensive decrease in intensity of the gravitational field with two separate epicenters caused by large granitoid massifs. Difference in the values of the field Δg is -50 mGal in marginal parts and is up to -68-76 mGal in epicenters. Both massifs are the marginal eastern part of a vast zone of magmatism, stretching in a submeridional direction far to the north beyond the studied area. Individual granitoid protrusions are clearly outlined on transformed maps Δg ($H_{\text{down}}=500$ m) and isolines V_{zz} in the form of local anomalies of a negative sign.

A negative gravitational anomaly is confined to the nuclear part of the Zhezkazgan syncline, within which two large local minimums (Northern and Southern) are fixed, the geological nature of which, despite the close parameters (area and intensity), is rather different. According to density selection of the Δg curves along the interpretation profiles, the northern minimum is apparently associated with the presence here of a thick layer of low-density deposits of the Zhidelisai suite, and the southern one, according to the interpretation, outlines the granitoids of the Terekty intrusive complex of the Middle Devonian, not exposed by erosion, at a depth of about 2900 meters and deeper. In addition, in the western marginal part of the southern minimum, a well has uncovered loose deposits of the Zhidelisai suite with deposits of rock salt, which can explain its nature. However, it is possible that both minimums are conditioned by the combined influence of the factors.



Figure 2. Transformations of the gravitational field. The isoline map V_{zz} :
 1 - Ulytau-Arganaty SFZ; 2 - Sarysu-Teniz SFZ; 3 - Konsky SFZ.

In the south of the site, south of the South-Terekty fault, the northern end of a large negative gravitational anomaly, according to the interpretation, also fixes the granitoids of the Terekty intrusive complex, not exposed by erosion, at a depth of about 3000 meters, composing a large massif of Talap.

To the west of the Zhezkazgan syncline, within the Zhanai and It-Auz brachyanticlines, a submeridionally elongated zone of a positive gravitational field was traced. This zone is associated with the approach to daylight surface of anticlinal structures anticipated at the depth of dense amphibolite-shale rocks of the Lower Proterozoic.

A number of local positive gravitational anomalies were distinguished in the

side parts of the Zhezkazgan syncline, the largest of which are the Spassky and South Sary-Oba maximums, within which respectively the Zhezkazgan deposit and the southern immersion of the Sary-Oba deposit are located. Their nature is presumably conditioned by an increase in density of ore horizons, mainly, epigenetically altered sedimentary host rocks.

The southern side of the Zhezkazgan syncline in the gravitational field has a clear band of positive gravi-maximums with a change in the strike from the sublatitudinal (Spassky maximum) to the northeast. A group of deposits and manifestations are confined to the gradient zones of this band: Spassky, Zhezkazgan, Ak-Chii, Dalny, Karashoshak, Sorkuduk, Zhartas, Pektas and Mezhdurechnoye. It should be noted that the nature of the most extensive and intensive Spassky gravi-maximum is presumably associated with the processes of intensive ore formation.

The Zhezkazgan copper deposit is structurally located on the western submerged lock part of the Kingir brachyanticline, at the junction of the Zhezkazgan-Terekty and the East Zhilandy deep faults. In this area, a fragment of the latter, according to regional works, is called East Ulytau.

The northern side of the Zhezkazgan syncline is accompanied by the Sary-Oba gravimaximum, to the gradient part of which confined are deposits of It-Auz and both Eastern and Western Sary-Oba. The entire northern group of deposits and manifestations, together with the above-mentioned Kipshakpai and Karashoshak, is confined to the junction of faults zones of the sublatitudinal strike (Kipshakpai throw) and a series of faults of the submeridional direction (Spassky, Eastern Zhilandy, Sary-Oba and a number of unnamed faults parallel to them), which are identified by gravimetric and, to a greater extent, seismic data.

In the studies of the geological structure of the Zhezkazgan ore region, utilised were maps of anomalous magnetic fields scaled 1: 100,000-1: 50,000, obtained by production companies upon high-precision aeromagnetic and ground surveys. Survey materials of different years were harmonised and interlinked. The measured magnetic field reflects various deep sources on the Earth's surface. Maps of the observed magnetic field provided the information about material composition of hidden Paleozoic structures and were utilized in specifying the outlining boundaries.

Most of existing methods of solving the inverse problem of magnetic exploration were developed for local anomalies. Therefore, in magnetometers, as in the analysis of the gravitational field, for practical purposes, the whole of geomagnetic field is divided into a normal field and an anomalous one. For this purpose, magnetic field transformations are calculated in order to emphasize certain features of a field and to weaken its side effects. As a rule, the anomalous

part is associated with the nonhomogeneous magnetization of Upper Paleozoic structures, and the normal part is associated with the effects of currents flowing in the Lower Paleozoic complexes and the Proterozoic crystalline basement (Istekova S.A. 2006).




Figure 3. The isoline map of the anomalous magnetic field ΔT_a
1 - Ulytau-Arganaty SFZ; 2 - Sarysu-Teniz SFZ; 3 - Konsky SFZ.

The magnetic field of the work area (figure 3) has a quiet character of intensity (ΔT)_a, varying from 100 to 50nTl, reflecting, in general, a non-magnetic carbonate-terrigenous-sedimentary thickness starting from the Middle Ordovician to the Upper Devonian-Permian ($\chi_{sr.} = 4-23 \cdot 10^{-5}$ units SI). Only a chain of positive magnetic anomalies (ΔT)_a, which in the north outside the reporting area ends with an intense magnetic anomaly (ΔT)_a associated with the manifestation of ultrabasites of the Esculy dome and accompanies the East

Ulytau fault, traced by fragments of Δg gradient zones of the northwest strike.

Small anomalies (ΔT)_a (over 300nTl) outline fragments of melanged mafic-ultramafic tectonic covers at depth. This increase in the magnetic field is associated with the juncture of two systems of discontinuous disturbances of the submeridional and sublatitudinal directions. In addition, positive magnetic field anomalies (ΔT)_a with a strength of over 50nTl are registered by deep, undiscovered, granodiorites of the Karamendy intrusive complex of the Early Devonian ($\chi_{sr.} = 636 \cdot 10^{-5}$ SI units).

The 1975-1977 Zhezkazgan geological exploration team carried out the seismic exploration work by reflection method within the Zhezkazgan ore region (Kotlyarov A.M., Smirnova N.N., etc.). The works clarified the structure of individual parts of the Zhezkazgan syncline, primarily the previously identified Sary-Oba anticline and the Kokdombak uplift. The reference seismic horizons were traced, and depths of productive deposits in the eastern and central parts of the Zhezkazgan syncline were determined (figure 4).




Figure 4. The geological and geophysical section along the III-III profile.

In 1980-84, experimental and methodological work was carried out on the site of the Northern Group of the Zhezkazgan type copper deposits, in order to evaluate the possibility of seismic exploration when searching directly for copper deposits by a number of anomalous signs (Smirnova N.N. et al., Zhezkazgan GEE). The authors analyzed the data, obtained by the past reflection method and the correlation refraction method works, in light of new velocity parameters for the Zhezkazgan syncline, and linked them with new data, obtained in the 1980-1981 MOGT survey.

In 1996-1997, the JSC “Zhezkazgan Geologiya” led by N.N. Smirnova conducted the MOGT seismic exploration by 24-fold profiling with exploratory drilling in the northeastern part of the Zhezkazgan Depression. The aim of the work was to identify the structures promising for cupreous sandstones, study the geological section of Upper Paleozoic deposits, trace tectonic disturbances and check the copper content of the identified structures. As a result, the team traced the reference reflecting horizons R_{III} (the sole of sediments C_1), R_{II} (the sole of sediments C_2 ts), R_{II} (the bottoms of the sole C_1v_2), the Terekty fault zone and other tectonic disturbances, and estimated the depth of the sole of the productive Taskuduk suite.

The research outcome. As a result of geological interpretation of the obtained geophysical materials, the geological structure of the Zhezkazgan ore district was clarified.

The gravitational and magnetic fields allowed us to study the heterogeneity of the deep geological structure (tectonics, intrusive magmatism). The qualitative analysis and results of the quantitative interpretation performed by both previous and present researchers allowed to identify the supposed contours of intrusions at depth: granitoids (according to gravity exploration), granodiorites and ultramafites (according to magnetic exploration), and to build a generalized structural and tectonic model of the district. According to the data of large-scale geophysical surveys, identified were the areas of development of presumably Zhidelisai deposits of considerable thickness (with lowered values of density $\sigma_{sr} = 2.60 \text{ g/cm}^3$ and of resistances 15-20 Ohm-m), as well as the areas of development of rocks of presumably Zhezkazgan and Kingir suites of considerable thickness (local gravimaximums, $\sigma_{sr} = 2.66-2.67 \text{ g/cm}^3$).

Tectonic disturbances were traced in space and at depth. Deep faults are usually accompanied by extended linearly elongated zones of high gradients of physical fields, regions of flank junction of anomalous zones of various strikes, clear-cut local anomalies of gravitational and magnetic fields - the local gravity minimums above the rock crush zones and the chains of positive anomalies (ΔT)_a, resulted from small intrusions of hyperbasites into weakened zones. The lithological-stratigraphic model of the district does not manifest itself in gravitational and magnetic fields, since sedimentary complexes of various formations do not differentiate by physical properties.

The gravitational and magnetic fields of the Zhezkazgan syncline district, in general, are differentiated ones. Apart from the geological structures mapped on daylight surface, the deep structure of the district and their features influence on the gravitational and magnetic fields. Within the territory under consideration, the southern locking of the Zhezkazgan syncline is accompanied by a quiet negative magnetic field (ΔT)_a, and is fixed by a local negative anomaly Δg , emphasizing the submeridional extension of the structure. The structure of the gravimetric field of the southern part of the Zhezkazgan SFZ is overwhelmingly influenced by the spread of a huge mass of Middle Paleozoic granitoids at depth, which cause the presence of local gravi-minimums.

In the southern part of the Zhezkazgan syncline, complicating folds of the second order, or the so-called structural “noses”, are located at the pass to the western periclinal locking of the Kingir brachyanticline: the Pokrovskaya and Krestovskaya anticlines and separating them the Zlatoust syncline, to which the Spassky gravi-maximum of sub-isometric shape (10×17 km) is confined with intensity of up to 3 mGal, coinciding with the Zhezkazgan deposit. This

gravi-maximum is conditioned by the denser composition of the underlying foundation as well as the higher density of epigenetically altered rocks of the host ore horizons (Strutynsky, 1988).

The Zhilandy syncline branches off from the northwestern end of the Zhezkazgan syncline and stretches in a meridional direction for almost 20 km. This narrow (3 to 5-6 km) continuous fold has the features of a typical linear fold with a rather complex internal structure. It splits into a number of synclinal pits, the southernmost of which is composed of deposits of the Taskuduk, Zhezkazgan and Zhidelisai suites. The undulating hinge of the syncline plunges in the direction of the Zhezkazgan syncline. The generalized lithological and stratigraphic horizons were identified as a result of the geological interpretation of the reflection method seismic data. According to the seismic data, fault zones are confidently identified by the shift of amplitudes of varying significance and the change in nature of the wave field (Martyshko P.S. et.al, 2016). The internal structure of the Zhezkazgan syncline is complicated by fault zones with different displacement amplitudes, and the sole depths isolines of the Famennian layer sediments emphasize the mosaic structure and the submeridional strike of the structure (figure 5).

Discussion. The deep structure of the Zhezkazgan syncline on sides of the Sary-Oba rise differs sharply, fixing the simple structure of the eastern part and the complex folded one of the western part. The deposits of the Zhezkazgan ($C_{2,3}$ đž) and overlying Zhidelisai (P_1 žd) and Kingir ($P_{1,2}$ kn) suites represent a conforming continuous section (in places, there are local disconformities over faults with repeated tectonic displacements), lying on sediments of the Taskuduk suite. Their accumulation coincides with the Middle Carboniferous tectonic activity, which led to formation of the Sarysu-Teniz uplift and division of the Zhezkazgan-Teniz coal basin into three independent depressions, Zhezkazgan, Shubarkul and Teniz (Boris S. 2010).

Application of modern means of registration, processing and interpretation of geological and geophysical data led to increased effectiveness of ore geophysics in studies of tectonic structure of ore areas, localization and morphology of differentiated intrusive massifs at depth, detection and deep mapping of ore-controlling faults, isolation of ore-controlling structures in sedimentary and effusive-sedimentary folded complexes, and determination of the spatial position of ore-bearing zones.




Figure 5. Geological and geophysical sections along the latitudinal and meridional profiles of the Zhezkazgan ore field.

Conclusion. Taking into account the rather complex geological structure of the given ore district and the significant differentiation of the composing rocks, ores and near-ore hosting strata, further identification of additional criteria for localization of copper mineralization is possible with the integration of detailed geological and geophysical methods, geochemical and physical-chemical studies of rocks and ores.

Information about the authors:

Istekova S.A. – Doctor of Geologo-Mineralogical Sciences, professor at the department of Geophysics Satbayev University Almaty, Kazakhstan; istekovy@mail.ru; <https://orcid.org/0000-0003-4298-7598>;

Aidarbekov Zh.K. – Field Engineer at Baker Hughes, PhD candidate at the Department of Geophysics Satbayev University Almaty, Kazakhstan, zh.aidarbekov@gmail.com; <https://orcid.org/my-orcid?orcid=0000-0002-4461-5256>.

REFERENCES

- [1] Aidarbekov Zh., Istekova S., Glass H. (2021) Complex of Geophysical Research for Studying Geological Structure of Zhezkazgan Ore Region in Kazakhstan. Conference Paper European Association of Geoscientists & Engineers. *Conference Proceedings, Engineering and Mining Geophysics 2021*, Gelendzhik Russia. 2021:1–9. ISSN 2658-6924. <https://doi.org/10.3997/2214-4609.202152070> (in Russ.).
- [2] Baibatsha A., Arshamov Y., Bekbotayeva A., Baratov R. (2017) Geology of the main

industrial types of copper ore deposits in Kazakhstan. International Multidisciplinary Scientific Geo Conference Surveying Geology and Mining Ecology Management, 11: 231-238. ISBN: 978-619-7105-98-8 ISSN: 1314-2704 <https://doi:10.5593/sgem2017/11/s01.029> (in Eng.).

[3] Balk P.I., Dolgal A.S., Balk T.V., Khristenko L.A. (2016) Minimax approach to inverse problems of geophysics, *Izvestiya Physics of the Solid Earth*, 52 -2: 242–253 <https://doi.org/10.1134/S1069351316020026> (in Eng.).

[4] Balk P.I., Dolgal A.S. (2017) New possibilities of increasing the informativeness of quantitative interpretation of gravity anomalies. *Doklady Earth Sciences*, 476-2:1168–1172. <https://doi.org/10.1134/S1028334X17100038> (in Eng.).

[5] Boris S., Stephen E. Box, John C. Wallis, (2010) Spatial Databases of Geological, Geophysical, and Mineral Resource Data Relevant to Sandstone-Hosted Copper Deposits in Central Kazakhstan: U.S. Geological Survey Scientific Investigations Report. Open-File Report: 1124, <https://doi.org/10.3133/ofr20101124> (in Eng.).

[6] Cossette P.M., Bookstrom A.A., Hayes T.S., Robinson G.R., Jr., Wallis J.C., Zientek M.L. (2014). Sandstone copper assessment of the Teniz Basin, Kazakhstan: U.S. Geological Survey Scientific Investigations Report, 2010–5090–R: 42 p. <http://dx.doi.org/10.3133/sir20105090R> (in Eng.).

[7] Hitzman M.W., Selley D. and Bull S. (2010) Formation of sedimentary rock-hosted stratiform copper deposits through Earth history: *Economic Geology*, 105-3: 627–639. <https://doi:10.2113/gsecongeo.105.3.627> (in Eng.).

[8] Istekova S.A. (2006) Modern methods of geophysical data analysis and interpretation [Sovremennyye metody analiza i interpretatsii geofizicheskikh dannykh]. *Izvestiya NAN RK. Seriya geologicheskaya*, 4:23-30. (in Russ.).

[9] Lyubetskaya L.D., Urdabayev A.T., Shabalina L.V., Bikeyev V.S. (2009) On the issue of the formation model of the Zhezkazgan deposit of medical sandstones. Conference on Satpayevskie chteniya, Almaty, Kazakhstan. (in Russ.).

[10] Martyshko P.S., Ladovsky I.V., Fedorova N.V., Byzov D.D., Tsidaev A.G. (2016) Theory and methods of complex combination of geophysical data, Yekaterinburg: Ural Branch of the Russian Academy of Sciences, 16 – 94 ISBN: 978-57691-2463-1 (in Russ.).

[11] Stephen E. Box., Reimar Seltmann, Michael L. Zientek, Boris Syusyura, Robert A. Creaser, Alla Dolgoplova. (2012) Dzhezkazgan and associated sandstone copper deposits of the Chu-Sarysu basin, Central Kazakhstan. *Society of Economic Geologists*, 303-328, <https://doi.org/10.3133/sir20105090E> (in Eng.).

[12] Togizov K.S., Zholtayev G.Z.; Isaeva L.D. (2019). The role of three-dimensional models of the deposit and thermodynamic conditions of its formation at selecting and evaluating resources of prospective sites, *News NAS RK. Series of geology and technical sciences*. 5(437):169-176. <https://doi:10.32014/2019.2518-170x.139> (in Eng.).

[13] Uzhkenov B.S. (2011) Innovationally technological schemes of forecasting and exploration jobs while exploring zhezkazgan typed ore fields. [Innovatsionnoye tekhnologicheskoye skhemy prognozno-poiskovykh rabot pri poiskakh mestorojdenii medi zhezkazganskogo tipa]. *Kazakhstan*, 3:68-71. (in Russ.).

[14] Zhezkazgan fields (2005) [Zhezkazganskoye mestorojdenie] Kazakhstan. ISBN: 9965-9746-3-2. (in Russ.).

[15] Zientek M.L., Chechetkin V.S., Parks H.L., Box S.E., Briggs D.A., Cossette P.M., Dolgoplova A., Hayes T.S., Seltmann R., Syusyura B., Taylor C.D., Wintzer N.E. (2014) Assessment of undiscovered sandstone copper deposits of the Kodar-Udokan area, Russia: U.S. Geological Survey Scientific Investigations Report. 2010–5090–M: 129. <http://dx.doi.org/10.3133/sir20105090M> (in Eng.).

CONTENTS

A.U. Abdullaev, Sh.S. Yusupov, L.Yu. Shin, A.V. Rasulov, Y.Zh. Yessenzhigitova HYDROGEOSEISMOLOGICAL PRECURSORS SUSAMYR EARTHQUAKE 1992.....	6
N.A. Abdimutalip, A.K. Kurbaniyazov, G. Toychibekova, G. Koishieva, G. Shalabaeva, N. Zholmagambetov INFLUENCE OF CHANGES IN THE LEVEL OF SALINITY OF THE ARAL SEA ON THE DEVELOPMENT OF ECOSYSTEMS.....	17
Zh.K. Aidarbekov, S.A. Istekova CLASSIFICATION OF GEOPHYSICAL FIELDS IN THE STUDY OF GEOLOGICAL AND STRUCTURAL FEATURES OF THE ZHEKAZGAN ORE DISTRICT.....	33
B. Almatova, B. Khamzina, A. Murzagaliyeva, A. Abdygalieva, A. Kalzhanova NATURAL SORBENTS AND SCIENTIFIC DESCRIPTION OF THEIR USE.....	49
Zh.A. Baimuratova, M.S. Kalmakhanova, SH.S.Shynazbekova, N.S. Kybyraeva, J.L. Diaz de Tuesta, H.T. Gomes MnFe ₂ O ₄ /ZHETISAY COMPOSITE AS A NOVEL MAGNETIC MATERIAL FOR ADSORPTION OF Ni(II).....	58
Ye.Z. Bukayev, G.K. Mutalibova, A.Z. Bukayeva A NEW TECHNOLOGY FOR MANUFACTURING POLYMER-CEMENT COMPOSITION FROM LIMESTONE-SHELL MINING WASTE.....	73
A.Zh. Kassenov, K.K. Abishev, A.S. Yanyushkin, D.A. Iskakova, B.N. Absadykov RESEARCH OF THE STRESS-STRAIN STATE OF HOLES WITH NEW BROACH DESIGNS.....	89
J.Kh. Khamroyev, K. Akmalaiuly, N. Fayzullayev MECHANICAL ACTIVATION OF NAVBAHORSK BENTONITE AND ITS TEXTURAL AND ADSORPTION CHARACTERISTICS.....	104

A.N. Kopobayeva, G.G. Blyalova, A. Bakyt, V.S. Portnov, A. Amangeldikyzy THE NATURE OF RARE EARTH ELEMENTS ACCUMULATION IN CLAY LAYERS AND COALS OF THE SHUBARKOL DEPOSIT.....	117
A. Leudanski, Y. Apimakh, A. Volnenko, D. Zhumadullayev, N. Seitkhanov CALCULATION OF FLOTATOR'S AERATOR FOR SEPARATION OF GROUND PLASTICS.....	131
Zh.T. Mukayev, M.M. Ulykpanova, Zh.O. Ozgeldinova, B.E. Kenzheshova, A.B. Khamitova CONTENT OF COPPER IN DESERT SOILS AND PLANTS OF EAST KAZAKHSTAN REGION.....	149
G. Sapinov, A. Imashev, Z. Mukhamedyarova CURRENT STATE OF THE PROBLEM OF MINING INDUCED SEISMICITY AND PROSPECT OF USING SEISMIC MONITORING SYSTEMS.....	161
V.G. Stepanets, V.L. Levin, G.K. Bekenova, M.S. Khakimzhanov, K.S. Togizov ACCESSORY COPPER ORE MINERALS AS A KEY ISSUE IN UNDERSTANDING THE GENESIS OF THE MAYATAS META-CARBONATITE ORES (ULYTAU, CENTRAL KAZAKHSTAN).....	172
S.A. Syedina, L.S. Shamganova, N.O. Berdinova, G.B. Abdikarimova MULTIVARIANT GEOMECHANICAL ESTIMATION OF THE DESIGN PARAMETERS' STABILITY OF SLOPE AND BENCH IN SOUTH SARBAI MINE.....	192
S.A. Tarikhazer, I.I. Mardanov INDICATORS OF ECOGEOMORPHOLOGICAL RISK FOR THE PURPOSE OF SUSTAINABLE DEVELOPMENT OF MOUNTAIN TERRITORIES.....	204
Zh.T. Tleuova, D.D. Snow, M.A. Mukhamedzhanov, E.Zh. Murtazin ASSESSMENT OF THE IMPACT OF HUMAN ACTIVITY ON GROUNDWATER STATUS OF SOUTH KAZAKHSTAN.....	217

Ye.A. Tseshkovskaya, A.T. Oralova, E.I. Golubeva, N.K. Tsoy, A.M. Zakharov	
DUST SUPPRESSION ON THE SURFACES OF STORAGE DEVICE OF TECHNOGENIC MINERAL FORMATIONS.....	230
B.T. Uakhitova, L.I. Ramatullaeva, M.K. Imangazin, M.M. Taizhigitova, R.U. Uakhitov	
ANALYSIS OF INJURIES AND PSYCHOLOGICAL RESEARCHES OF WORKERS IN THE MELTING SHOPS OF THE AKTUBINSK FERRALOYS PLANT.....	242
G.T. Shakulikova, S.M. Akhmetov, A.N. Medzhidova, N.M. Akhmetov, Zh.K. Zaidemova	
IMPROVING THE DESIGN OF INCLINED WELLS AS THE BASIS FOR THE DEVELOPMENT OF HARD-TO-RECOVER HYDROCARBON RESERVES.....	259
K.T. Sherov, M.R. Sikhimbayev, B.N. Absadykov, T.K. Balgabekov, A.D. Zhakaba	
STUDY OF TEMPERATURE DISTRIBUTION DURING ROTARY TURNING OF WEAR-RESISTANT CAST IRON.....	271

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://www.geolog-technical.kz/index.php/en/>

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Редакторы: *М.С. Ахметова, А. Ботанқызы, Д.С. Аленов, Р.Ж. Мрзабаева*

Верстка на компьютере *Г.Д.Жадыранова*

Подписано в печать 19.04.2022.

Формат 70x90^{1/16}. Бумага офсетная. Печать – ризограф.

11,5 п.л. Тираж 300. Заказ 2.