

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ГЕОЛОГИЯ ЖӘНЕ ТЕХНИКАЛЫҚ ҒЫЛЫМДАР
СЕРИЯСЫ



СЕРИЯ
ГЕОЛОГИИ И ТЕХНИЧЕСКИХ НАУК



SERIES
OF GEOLOGY AND TECHNICAL SCIENCES

2 (422)

НАУРЫЗ – СӘУІР 2017 ж.
МАРТ – АПРЕЛЬ 2017 г.
MARCH – APRIL 2017

ЖУРНАЛ 1940 ЖЫЛДАН ШЫҒА БАСТАҒАН
ЖУРНАЛ ИЗДАЕТСЯ С 1940 г.
THE JOURNAL WAS FOUNDED IN 1940.

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р ы

э. ғ. д., профессор, ҚР ҰҒА корреспондент-мүшесі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Р е д а к ц и я а л қ а с ы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., корр.-мүшесі (Қазақстан)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қырғыстан)
Беспәев Х.А. проф. (Қазақстан)
Бишимбаев В.К. проф., академик (Қазақстан)
Буктуков Н.С. проф., корр.-мүшесі (Қазақстан)
Бұлат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Ерғалиев Г.Х. проф., академик (Қазақстан)
Жуков Н.М. проф. (Қазақстан)
Кенжалиев Б.К. проф. (Қазақстан)
Қожахметов С.М. проф., академик (Қазақстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., корр.-мүшесі (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Нигматова С.А. проф. (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Қазақстан)
Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)
Сейтмуратова Э.Ю. проф., корр.-мүшесі (Қазақстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«ҚР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://nauka-nanrk.kz/geology-technical.kz>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р

д. э. н., профессор, член-корреспондент НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н н а я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., чл.-корр. (Казахстан)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Беспаяев Х.А. проф. (Казахстан)
Бишимбаев В.К. проф., академик (Казахстан)
Буктуков Н.С. проф., чл.-корр. (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Ергалиев Г.Х. проф., академик (Казахстан)
Жуков Н.М. проф. (Казахстан)
Кенжалиев Б.К. проф. (Казахстан)
Кожаметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., чл.-корр. (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Нигматова С.А. проф. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Казахстан)
Сейтов Н.С. проф., чл.-корр. (Казахстан)
Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

<http://наука-нанрк.kz/geology-technical.kz>

© Национальная академия наук Республики Казахстан, 2017

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of Economics, professor, corresponding member of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

E d i t o r i a l b o a r d:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., corr. member. (Kazakhstan)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Bespayev Kh.A. prof. (Kazakhstan)
Bishimbayev V.K. prof., academician (Kazakhstan)
Buktukov N.S. prof., corr. member. (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadjikistan)
Gravis R.M. prof. (USA)
Yergaliev G.Kh. prof., academician (Kazakhstan)
Zhukov N.M. prof. (Kazakhstan)
Kenzhaliyev B.K. prof. (Kazakhstan)
Kozhakhmetov S.M. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., corr. member. (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Nigmatova S.A. prof. (Kazakhstan)
Ozdoyev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Rakishev B.R. prof., academician (Kazakhstan)
Seitov N.S. prof., corr. member. (Kazakhstan)
Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 2, Number 422 (2017), 212 – 221

UDC 517.928.4:517.929.4

Zh. K. Dyusembina

Eurasian national university named after L. N. Gumilev, Astana, Kazakhstan.

E-mail: zdyusembina@mail.ru

MODIFICATION OF THE QUASILINEAR CONTROL SYSTEM OF BIOMEDICINE

Abstract. The stability of the respiratory chemostat on a finite time interval was studied and the task of stabilizing the biomedical system of the respiratory chemostat for linear and quasilinear control systems was solved. The study of such regulatory processes in the body plays an important role in the development of technical life support systems, and this applies not only to the voltage regulating system of the of carbon dioxide and oxygen (the "respiratory chemostat" system), but also to a number of other vital systems. First of all, this refers to the "cardiovascular chemostat" system, which, as a biological self-regulation system, has the task of washing the tissue reservoir with fresh blood at such a rate as to maintain the voltage of carbon dioxide and oxygen in tissues at normal or near levels. This is a complex hydrodynamic system of biological self-regulation, the numerous parameters of which are under continuous influence of various kinds of control signals, which in turn depend on external conditions relative to the organism and perturbations. The regulating regime of this system is continuously subjected to sharply and suddenly changing external influences, which have an extreme character.

Keywords: respiratory chemostat, control system, biological system, finite interval, stability.

The problem of the stability of the medical and biological system is one of the main problems of system analysis, mathematical ecology and biomedicine. The task of synthesizing vector control from a given class for a managed medical-biological system and population dynamics of a community with self-limiting by mathematical models is represented in the form of quasilinear equations. The models of biomedicine are considered and the quasi-linear control system of the respiratory chemostat on a finite time interval is investigated.

Changing $F_{CO_2}^I$, we act on an isolated controlled system, it behaves like a simple linear system with constant coefficients. Concentration of $CO_2(F_{CO_2}^I)$ functions as a direct driving force. When the respiratory center affects the controlled system, this effect is manifested through ventilation \dot{V}_A . When considering the behavior of the chemostat in the steady state, we assumed for convenience that the input signals of the control system are determined by the levels of $pCO_2, [H^+]$ and pO_2 in arterial blood [1].

Receptors are really located on the path of arterial blood, but proceed only from the fact that in each particular considered steady situation the values of these parameters are closely related to their effective values in those places where the real receptors are located.

Let us consider the time course of pulmonary ventilation and changes of pCO_2 in arterial blood with a sudden change in the concentration of carbon dioxide in the inspired air.

To simplify the analysis, we introduce a number of additional assumptions: 1) the lungs are a reservoir of constant volume vented by a continuous gas stream with a dead space equal to zero, and with a uniform composition; 2) the respiratory coefficient (RQ) at each moment of time is equal to one; 3) transport delays in the transfer of blood are negligible; 4) the respiratory center and other tissues are a homogeneous reservoir, washed by a constant flow of blood; 5) arterial blood, venous blood and "tissues"

are characterized by the same linearized absorption curve CO_2 ; 6) the partial pressures CO_2 in the exhaled air and in the alveolar air and the tension CO_2 in the arterial blood are always equal to each other, as well as the stresses CO_2 in the "tissues" and in the venous blood; 7) the control system is a simple inertial-free (that is, does not contain dynamic elements) proportional regulator.

To obtain equations describing the dynamic behavior of the biomedical system, we use the same general principles of equilibrium and continuity, on which Newton's Laws of motion and Kirchhoff's laws for electric circuits are based. We start with the pulmonary or alveolar reservoir and write down the "continuity equation for carbon dioxide", which states that the rate of change in concentration of CO_2 in the alveolar gas $\dot{\theta}_A$ is equal to the quotient of dividing the difference between the rates of intake and washing out of carbon dioxide by the volume of the reservoir K_A [1]:

$$\dot{\theta}_A = \frac{1}{K_A} [\dot{V}_A F_{\text{CO}_2}^I + q_3 - q_1 - q_2] \quad (1)$$

Carbon dioxide enters the lungs with inhaled gas at a rate equal to the product of pulmonary ventilation by the concentration of $F_{\text{CO}_2}^I$ in the inspired air, and with venous blood at the rate q_3 , and leaves the lungs with exhaled air at the rate q_1 and with arterial blood at the rate q_2 . Next, we will write down another equation of continuity for the tissue reservoir: the rate of change in the concentration of carbon dioxide in the tissues ($\dot{\theta}_T$) is equal to the sum of the rates of formation of CO_2 during the exchange (MR) of CO_2 with arterial blood flow (q_2) and leaching of CO_2 with venous blood (q_3) divided by the volume of the tissue reservoir K_T :

$$\dot{\theta}_T = \frac{1}{K_T} [MR + q_2 - q_3]. \quad (2)$$

Let us write down three equilibrium equations.

The first of them reflects the equality of concentrations of carbon dioxide in the alveolar and exhaled air:

$$\theta_A = \frac{q_2}{\dot{V}_A}. \quad (3)$$

The second equation describes the equilibrium of concentrations of CO_2 in the alveolar air and in the arterial blood, taking into account the linearized CO_2 absorption curve:

$$\frac{q_2}{Q} = BA_S(\theta_A) + A_i, \quad (4)$$

where Q minute volume of the heart, B - atmosphere pressure, A_S and A_i - respectively, the slope of the linear absorption curve and the ordinate of the point of its intersection with the ordinate axis.

The third equation expresses the equality of carbon dioxide concentrations in tissues and venous blood:

$$\theta_T = \frac{q_3}{Q}. \quad (5)$$

To study the behavior of an isolated controlled system, it is necessary to solve this system of equations. There are different ways for this, but in any case, we should, first of all, choose which of the five dependent variables is considered the output (or outputs) of the system and which of the nine independent variables ($\dot{V}_A, F_{\text{CO}_2}^I, MR, K_A, K_T, Q, B, A_S, A_i$) is considered to be the input (or inputs) of the system. We choose the variables θ_T and θ_A as the outputs of the system, and as the input - $F_{\text{CO}_2}^I(t)$. Thus, we assume that the variables $\dot{V}_A, MR, K_A, K_T, Q, B, A_S$ and A_i do not change with time. Combine the five obtained basic equations so as to get a differential equation with respect to one of the dependent variables, θ_T or θ_A . So, if we solve equation (3) relative to q_1 , (4) - relative to q_2 and equation (5) - relative to q_3 .

Substituting these values into equations (1) and (2), we obtain

$$\dot{\theta}_A = \frac{1}{K_A} [\dot{V}_A (F_{CO_2}^I - \theta_A) + Q(\theta_T - BA_S \theta_A - A_i)] \quad (6)$$

and

$$\dot{\theta}_A = \frac{1}{K_A} [MR - Q(\theta_T - BA_S \theta_A - A_i)]. \quad (7)$$

Solving equation (7) with respect to θ_A , we obtain

$$\theta_A = \frac{1}{QBA_S} (K_T \dot{\theta}_T - MR - QA_i + Q\theta_T). \quad (8)$$

The differentiation of (8) gives

$$\dot{\theta}_A = \frac{1}{QBA_S} (K_T \ddot{\theta}_T + Q\dot{\theta}_T). \quad (9)$$

Substituting (8) and (9) into (6), we obtain the required equation relative to θ_T :

$$\frac{K_A K_T}{Q \dot{V}_A} \ddot{\theta}_T + \left[\frac{K_A}{\dot{V}_A} + \frac{K_T BA_S}{\dot{V}_A} + \frac{K_T}{Q} \right] \dot{\theta}_T + \theta_T = BA_S F_{CO_2}^I(t) + \frac{BA_S MR}{\dot{V}_A} + \frac{MR}{Q} + A_i. \quad (10)$$

Having obtained the equation with respect to θ_T , we can find θ_A in several ways. From equation (8), find the expression for the concentration θ_A , substituting in (8) the expressions for θ_T and $\dot{\theta}_T$. But there is another method of determination θ_A , analogous to the method used to derive equation (10) with respect to θ_T . It consists in the fact that equation (6) is solved relatively to θ_T , the result obtained is differentiated (in this case, an expression for $\dot{\theta}_T$ is obtained) and the expressions for θ_T and $\dot{\theta}_T$ in equation (7) are substituted. The equation derived in this way has the following form [5]:

$$\frac{K_A K_T}{Q \dot{V}_A} \ddot{\theta}_A + \left[\frac{K_A}{\dot{V}_A} + \frac{K_T BA_S}{\dot{V}_A} + \frac{K_T}{Q} \right] \dot{\theta}_A + \theta_A = F_{CO_2}^I(t) + \frac{K_T}{Q} \dot{F}_{CO_2}^I(t) + \frac{MR}{\dot{V}_A}. \quad (11)$$

Equations (10) and (11) describe the behavior of an isolated controlled system under the assumption that the driving function is the concentration of carbon dioxide in the inspired air $F_{CO_2}^I(t)$. We solve these equations with respect to $\theta_T(t)$ and $\theta_A(t)$ for the case when $F_{CO_2}^I(t)$ represents a jump with amplitude $(F_{CO_2}^I(t))_1$. To solve these equations, we apply the classical method or the Laplace transform method.

The control system should be a "simple proportional regulator that does not contain dynamic elements". We suppose now that the same relation can be written in such a way that the concentration θ_T is included in it, and that it is valid for both transitional and steady-state conditions. We write down the equation of the control system in the following form:

$$\dot{V}_A = k_p [\theta_T - \theta_{Ti}] + \dot{V}_{Ar}. \quad (12)$$

In this equation θ_{Ti} - preset value, \dot{V}_{Ar} - bias signal. The gain of the proportional regulator is k_p .

Equations of motion (12) can be reduced to the following form, assuming $y = \theta_T$:

$$M \frac{d^2 y}{dt^2} + R \frac{dy}{dt} + Ky = F, \quad (13)$$

where

$$M = \frac{K_A K_T}{Q \dot{V}_A}, \quad R = \frac{K_A}{\dot{V}_A} + \frac{BA_S K_T}{\dot{V}_A} + \frac{K_T}{Q}, \quad K = 1, \quad F = BA_S F_{CO_2}^I(t) + \frac{BA_S MR}{\dot{V}_A} + \frac{MR}{Q} + A_i.$$

The angular frequency: $\varpi_n = \left(\frac{K}{M}\right)^{\frac{1}{2}} = \frac{1}{\sqrt{M}}$.

Attenuation factor: $b = \frac{R}{2(KM)^{\frac{1}{2}}} = \frac{R}{2\sqrt{M}}$.

Consequently, the equation (13) takes the form:

$$\frac{1}{\varpi_n^2} \cdot \frac{d^2 y}{dt^2} + \frac{2b}{\varpi_n} \cdot \frac{dy}{dt} + y = \left[\frac{1}{K}\right] F = F, \quad (14)$$

or

$$\frac{d^2 y}{dt^2} + 2b \cdot \varpi_n \frac{dy}{dt} + \varpi_n^2 y = \varpi_n^2 F \quad (15)$$

with boundary conditions

$$y(0) = y_0, \quad \dot{y}(0) = y_1, \quad y(T) = y_T, \quad \dot{y}(T) = 0.$$

Using the change of variables

$$x = y - y_T$$

we have that

$$\frac{d^2 x}{dt^2} + 2b \cdot \varpi_n \frac{dx}{dt} + \varpi_n^2 x = \varpi_n^2 (F - y_T) = u,$$

where

$$x = x_1, \quad u = \varpi_n^2 (F - y_T).$$

This differential equation in the Cauchy normal form takes the form:

$$\begin{cases} \dot{x}_1 = x_2 \\ \dot{x}_2 = -\varpi_n^2 x_1 - 2b \varpi_n x_2 + u \end{cases} \quad (16)$$

with boundary conditions

$$\begin{aligned} x_1(0) &= x_{10} = y_0 - y_T, \\ x_2(0) &= x_{20} = y_1, \\ x_1(T) &= 0, \\ x_2(T) &= 0. \end{aligned} \quad (17)$$

In the vector-matrix form:

$$\dot{x} = Ax + Bu, \quad x(0) = x_0, \quad x(T) = 0,$$

where

$$A = \begin{pmatrix} 0 & 1 \\ -\varpi_n^2 & -2b\varpi_n \end{pmatrix}, \quad B = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad u - \text{scalar}.$$

Fundamental system

$$\hat{O}(t) = A\hat{O}(t), \quad \hat{O}(0) = \hat{A}$$

is represented as:

$$\begin{pmatrix} \hat{O}_{11} & \hat{O}_{12} \\ \hat{O}_{21} & \hat{O}_{22} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\varpi_n^2 & -2b\varpi_n \end{pmatrix} \begin{pmatrix} \hat{O}_{11} & \hat{O}_{12} \\ \hat{O}_{21} & \hat{O}_{22} \end{pmatrix} = \begin{pmatrix} \hat{O}_{21} & \hat{O}_{22} \\ -\varpi_n^2 \hat{O}_{11} - 2b\varpi_n \hat{O}_{21} & -\varpi_n^2 \hat{O}_{12} - 2b\varpi_n \hat{O}_{22} \end{pmatrix}.$$

Hence we have

$$\begin{cases} \dot{\Phi}_{11} = \Phi_{21}, \\ \dot{\Phi}_{21} = -\varpi_n^2 \Phi_{11} - 2b\varpi_n \Phi_{21}, \\ \dot{\Phi}_{12} = \Phi_{22}, \\ \dot{\Phi}_{22} = -\varpi_n^2 \Phi_{12} - 2b\varpi_n \Phi_{21} \Phi_{22}, \end{cases} \quad (18)$$

$$\hat{O}_{11}(0) = 1, \quad \hat{O}_{21}(0) = 0, \quad \hat{O}_{12}(0) = 0, \quad \hat{O}_{22}(0) = 1.$$

Next we get

$$\ddot{O}_{11} + 2b\omega_n \dot{O}_{11} + \omega_o^2 \hat{O}_{11} = 0, \quad \ddot{O}_{12} + 2b\omega_n \dot{O}_{12} + \omega_o^2 \hat{O}_{12} = 0. \quad (19)$$

We note that the disturbance $F_{CO_2}^I(s)$ is the direct driving force for the controlled system, and the control value $\dot{V}_A(s)$ excites the system due to parametric influence. The latter is the source of the nonlinearity of the system, as can be verified by substituting the equation (6) in (19).

We obtain the following equation of a closed system with respect to θ_T :

$$a\ddot{\theta}_T + \beta\dot{\theta}_T + \gamma\theta_T\theta_T + \theta_T^2 + \eta\theta_T = \lambda, \quad (20)$$

where

$$\begin{aligned} \alpha &\equiv \frac{K_A K_T}{k_p Q}, \\ \beta &\equiv \frac{K_A + K_T B A_S}{k_p} - \frac{K_T \theta_{Ti}}{Q} + \frac{K_T \dot{V}_{Ar}}{k_p Q}, \quad \gamma \equiv \frac{K_T}{Q}, \\ \eta &\equiv \frac{\dot{V}_{Ar}}{k_p} - \theta_{Ti} - B A_S F_{CO_2}^I - \frac{MR}{Q} - A_i, \\ \lambda &= \frac{B A_S MR}{k_p} + \left[\frac{\dot{V}_{Ar}}{k_p} - \theta_{Ti} \right] \left[B A_S F_{CO_2}^I + \frac{MR}{Q} + A_i \right]. \end{aligned}$$

The equation (20) is a nonlinear differential equation, since it contains terms of the second degree $\dot{\theta}_T \theta_T$ and θ_T^2 . Since \dot{V}_A is a linear algebraic function of θ_T concentration (equation 6), then the equation of a closed system with respect to \dot{V}_A in the form would be identical to equation (20). Relatively to θ_A , a complex nonlinear equation can also be obtained [2].

Let us consider the stabilization of the motions of the quadratic system (20) on a finite interval of time [4].

We denote $\bar{u} = \dot{V}_{Ar}$ – as control

$$\begin{aligned} \beta &= \beta_1 + \beta_2 \bar{u}, \\ \eta &= \eta_1 \bar{u} - \eta_2, & \gamma_1 &= \gamma / \alpha, & \lambda_1 &= \lambda / \alpha, & \eta_1 &= 1 / \alpha, \\ \lambda &= \lambda_1 + \lambda_2 \bar{u}, \end{aligned}$$

where

$$\beta_1 = \frac{\left[\frac{K_A + K_T BA_S}{k_p} - \frac{K_T Q_{T_i}}{Q} \right]}{\alpha}, \quad \beta_2 = \frac{K_T}{k_p Q^\alpha},$$

$$\eta_1 = \frac{1}{K_p \alpha}, \quad \eta_2 = \frac{\left[Q_{T_i} - BA_S F_{co}^I - \frac{MR}{Q} - A_i \right]}{\alpha},$$

$$\lambda_1 = \frac{\left[\frac{BA_S MR}{k_p} - Q_{T_i} \left(BA_S F_{co}^I + \frac{MR}{Q} + A_i \right) \right]}{\alpha},$$

$$\lambda_2 = \frac{1}{k_p \alpha} \left(BA_S F_{co}^I + \frac{MR}{Q} + A_i \right).$$

Then,

$$\ddot{\theta}_T + \beta_1 \dot{\theta}_T - \eta_2 \theta_T + \beta_2 \bar{u} \dot{\theta}_T + \gamma_1 \dot{\theta}_T \theta_T + \eta_2 \theta_T^2 + \eta_1 \bar{u} \theta_T = \lambda_1 + \lambda_2 \bar{u}. \tag{21}$$

We consider,

$$\theta_T = y,$$

$$\dot{y} + \beta_1 \dot{y} - \eta_2 y + \beta_2 \bar{u} \dot{y} + \gamma_1 \dot{y} y + \eta_1 y^2 + \eta_1 \bar{u} y = \lambda_1 + \lambda_2 \bar{u}.$$

$$y(0) = y_0, \quad \dot{y}(0) = y_1, \quad y(T) = y_T, \quad \dot{y}(T) = 0.$$

Then,

$$x = y - y_T,$$

$$\ddot{x} + \beta_1 \dot{x} - \eta_2 (x + y_T) + \beta_2 \bar{u} \dot{x} + \gamma_1 \dot{x} (x + y_T) + \eta_1 (x + y_T)^2 + \eta_1 \bar{u} (x + y_T) = \lambda_1 + \lambda_2 \bar{u}$$

or

$$\ddot{x} + \beta_1 \dot{x} + \eta_3 x + \beta_2 \bar{u} \dot{x} + \gamma_1 \dot{x} x + \eta_2 x^2 + \eta_1 \bar{u} x + \mu_1 = \lambda_3 \bar{u}, \tag{22}$$

where

$$\mu_1 = -\eta_2 y_T + \gamma_1 y_T + \eta_1 y_T^2 - \lambda_1,$$

$$\lambda_3 = \lambda_2 - \eta_1 y_T, \quad \eta_3 = 2\eta_1 y_T - \eta_2,$$

$$x(0) = y_0 - y_T, \quad \dot{x}(0) = y_1,$$

$$x(T) = y_T - y_T = 0, \quad \dot{x}(T) = \dot{y}(T) = 0.$$

We consider,

$$u = \lambda_3 \bar{u} - \mu_1, \quad \bar{u} = \frac{1}{\lambda_3} (u + \mu_1), \quad \ddot{x} + \beta_4 \dot{x} + \eta_5 x + \beta_3 u \dot{x} + \gamma_1 \dot{x} x + \eta_1 x^2 + \eta_4 u x = u, \tag{23}$$

where

$$\beta_3 = \frac{\beta_2}{\lambda_3}, \quad \beta_4 = \beta_1 + \frac{\mu_1}{\lambda_3}, \quad \eta_4 = \frac{\eta_1}{\lambda_3}, \quad \eta_5 = \eta_3 + \frac{\eta_1 \mu_1}{\lambda_3}.$$

Let it be

$$\begin{cases} x = x_1, \\ \dot{x}_1 = x_2, \\ \dot{x}_2 = -\eta_5 x_1 - \beta_4 x_2 + u + f_2(x, u), \end{cases} \tag{24}$$

$$\begin{aligned} f_2(x, u) &= -\beta_3 u x_2 - \gamma_1 x_1 x_2 - \eta_1 x_1^2 - \eta_4 x_1 u, & t \in [0, T] \\ x_1(0) &= x_{10}, & x_1(T) &= 0, \\ x_2(0) &= x_{20} = y_1, & x_2(T) &= 0. \end{aligned}$$

We consider a linear system

$$\dot{x} = Ax + Bu^0.$$

Fundamental system

$$\hat{O}(t) = A\hat{O}(t), \quad \hat{O}(0) = E$$

has the form:

$$\begin{pmatrix} \hat{O}_{11} & \hat{O}_{12} \\ \hat{O}_{21} & \hat{O}_{22} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\beta_4 & -\eta_5 \end{pmatrix} \begin{pmatrix} \hat{O}_{11} & \hat{O}_{12} \\ \hat{O}_{21} & \hat{O}_{22} \end{pmatrix} = \begin{pmatrix} \hat{O}_{21} & \hat{O}_{22} \\ -\beta_4 \hat{O}_{11} - \eta_5 \hat{O}_{21} & -\beta_4 \hat{O}_{12} - \eta_5 \hat{O}_{22} \end{pmatrix}.$$

At that

$$\begin{aligned} \dot{\Phi}_{11} &= \Phi_{21}, \\ \dot{\Phi}_{21} &= -\beta_4 \Phi_{11} - \eta_5 \Phi_{21}, \\ \dot{\Phi}_{12} &= \Phi_{22}, \\ \dot{\Phi}_{22} &= -\beta_4 \Phi_{12} - \eta_5 \Phi_{22}, \\ \hat{O}_{11}(0) &= 1, \quad \hat{O}_{12}(0) = 0, \quad \hat{O}_{21}(0) = 0, \quad \hat{O}_{22}(0) = 1, \\ \bar{O}_{21} + \eta_5 \hat{O}_{21} + \beta_4 \hat{O}_{21} &= 0, & (25) \\ \bar{O}_{22} + \eta_5 \hat{O}_{22} + \beta_4 \hat{O}_{22} &= 0. & (26) \end{aligned}$$

The characteristic equation:

$$\lambda^2 + \eta_5 \lambda + \beta_4 = 0, \quad \lambda_{1,2} = \frac{-\eta_5 \pm \sqrt{\eta_5^2 - 4\beta_4}}{2}.$$

has the roots

$$\begin{aligned} \lambda_1 &= -\mu_1, \quad \mu_1 = \frac{1}{2}[\eta_5 - \sqrt{D}] > 0, \\ \lambda_2 &= -\mu_2, \quad \mu_2 = \frac{1}{2}[\eta_5 + \sqrt{D}] > 0. \end{aligned}$$

The fundamental matrix of solutions has the form:

$$\Phi(t) = \frac{1}{\mu_2 - \mu_1} \begin{pmatrix} [(\mu_2 - \eta_5)e^{-\mu_2 t} + (\eta_5 - \mu_1)e^{-\mu_1 t}] & \frac{-(\mu_2 - \eta_5)(\mu_1 - \eta_5)}{\beta_4} (e^{-\mu_2 t} - e^{-\mu_1 t}) \\ \beta_4 (e^{-\mu_2 t} - e^{-\mu_1 t}) & [(\mu_2 - \eta_5)e^{-\mu_1 t} - (\mu_1 - \eta_5)e^{-\mu_2 t}] \end{pmatrix} = \frac{1}{\mu_2 - \mu_1} \bar{\Phi}(t).$$

The inverse matrix:

$$\begin{aligned} \hat{O}^{-1}(t) &= (\mu_2 - \mu_1) \bar{O}^{-1}(t), \\ \Delta &= e^{-(\mu_1 + \mu_2)t} [(\mu_2 - \eta_5) - (\mu_1 - \eta_5)]^2 = e^{-(\mu_1 + \mu_2)t} (\mu_2 - \mu_1)^2 > 0. \end{aligned}$$

Consequently,

$$\hat{O}^{-1}(t) = \frac{\mu_2 - \mu_1}{\Delta} \begin{pmatrix} (\mu_2 - \eta_5)e^{-\mu_1 t} - (\mu_1 - \eta_5)e^{-\mu_2 t}, & \frac{(\mu_2 - \eta_5)(\mu_1 - \eta_5)}{\beta_4} (e^{-\mu_2 t} - e^{-\mu_1 t}) \\ -\beta_4 (e^{-\mu_2 t} - e^{-\mu_1 t}), & (\mu_2 - \eta_5)e^{-\mu_2 t} + (\eta_5 - \mu_1)e^{-\mu_1 t} \end{pmatrix}.$$

we get

$$Q(t) = \hat{O}^{-1}(t) B = \frac{(\mu_2 - \mu_1)}{\Delta} \begin{pmatrix} \frac{(\mu_2 - \eta_5)(\mu_1 - \eta_5)}{\beta_4} (e^{-\mu_2 t} - e^{-\mu_1 t}) \\ (\mu_2 - \eta_5)e^{-\mu_2 t} + (\eta_5 - \mu_1)e^{-\mu_1 t} \end{pmatrix},$$

and

$$Q(t)Q^{\alpha}(t) = \frac{(\mu_2 - \mu_1)^2}{\Delta^2} \begin{pmatrix} \Phi_{11}(t), & \Phi_{12}(t) \\ \Phi_{12}(t), & \Phi_{22}(t) \end{pmatrix},$$

where

$$\begin{aligned} \hat{O}_{11}(t) &= \frac{(\mu_2 - \eta_5)^2(\mu_1 - \eta_5)^2}{\beta_4^2} (e^{-\mu_2 t} - e^{-\mu_1 t})^2, \\ \hat{O}_{12}(t) &= [(\mu_2 - \eta_5)e^{-\mu_2 t} + (\eta_5 - \mu_1)e^{-\mu_1 t}] \frac{(\mu_2 - \eta_5)(\mu_2 - \eta_5)}{\beta_4} (e^{-\mu_2 t} - e^{-\mu_1 t}), \\ \hat{O}_{22}(t) &= [(\mu_2 - \eta_5)e^{-\mu_2 t} + (\eta_5 - \mu_1)e^{-\mu_1 t}]^2, \\ R(t, T) &= \int_t^T Q(\tau)Q^*(\tau)d\tau = \begin{pmatrix} R_{11}, & R_{12} \\ R_{12}, & R_{22} \end{pmatrix} \frac{(\mu_2 - \mu_1)^2}{\Delta^2}, \\ R_{11}(t) &= \int_t^T \Phi_{11}(\tau)d\tau, \quad R_{12}(t) = \int_t^T \Phi_{12}(\tau)d\tau, \quad R_{22}(t) = \int_t^T \Phi_{22}(\tau)d\tau. \end{aligned}$$

Let us check:

$$R(o, T) > 0, \text{ т.е. } R_{11}(0) > 0, \quad R_{11}(0)R_{22}(0) - R_{12}^2(0) > 0, \text{ using } T > 0.$$

We calculate $R^{-1}(0, T)$:

$$\begin{aligned} W(t, T) &= \hat{O}(t) R(t, T) \hat{O}^*(t) = \begin{pmatrix} W_{11} & W_{12} \\ W_{21} & W_{22} \end{pmatrix}, \\ K(t) &= W^{-1}(t, T) = \begin{pmatrix} K_{11}(t), & K_{12}(t) \\ K_{12}(t), & K_{22}(t) \end{pmatrix}. \end{aligned}$$

Then the stabilizing control will take the form:

$$u^0(t, x) = -B^* K(t)x = -(0, 1) \begin{pmatrix} K_{11}(t), & K_{12}(t) \\ K_{12}(t), & K_{22}(t) \end{pmatrix} x = -K_{12}(t)x_1 - K_{22}(t)x_2.$$

$$u(t, x) = u^0(t, x) + v(t, x), \quad t \in [0, T],$$

where the scalar function $v(t, x)$ is determined under the condition [3]

$$2B v(t, x) + 2f(-B^* K(t)x + v(t, x), x) - BB^* K(t)x = -\bar{R}(t)x, \tag{27}$$

where $\bar{R}(t)$ – 2×2 matrix such that

$$K\bar{R}(t) + \bar{R}^*(t)K \geq 0,$$

ie, non-negative definite.

Let us suppose that

$$\begin{aligned} \bar{R}_{11} &= 0, \quad \bar{R}_{12} = 0, \\ \bar{R} &= \begin{pmatrix} 0 & 0 \\ \bar{R}_{21}(t) & \bar{R}_{22}(t) \end{pmatrix}. \end{aligned}$$

From (27) we have the following equation for determining $v(t, x)$.

$$\begin{aligned} 2 v(t, x) - 2\{\gamma_1 x_1 x_2 + \eta_1 x_1^2 + (\beta_3 x_2 + \eta_4 x_1)(-K_{12} x_1 - K_{22} x_2)\} - \\ - 2(\beta_3 x_2 + \eta_4 x_1)v(t, x) - K_{12} x_1 - K_{22} x_2 = -\bar{R}_{21} x_1 - \bar{R}_{22}(t) x_2. \end{aligned} \quad (28)$$

Let it be

$$\begin{aligned} \bar{R}_{21} &= K_{12}, \quad \bar{R}_{22} = K_{22}, \\ K\bar{R} &= \begin{pmatrix} K_{11} & K_{12} \\ K_{12} & K_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ K_{12} & K_{22} \end{pmatrix} = \begin{pmatrix} K_{12}^2 & K_{12}K_{22} \\ K_{12}K_{22} & K_{22}^2 \end{pmatrix}, \\ K\bar{R} + \bar{R}^* K &= 2 \begin{pmatrix} K_{12}^2 & K_{12}K_{22} \\ K_{12}K_{22} & K_{22}^2 \end{pmatrix} \geq 0, \quad K_{12}^2 > 0, \\ K_{12}^2 K_{22}^2 - K_{12}^2 K_{22}^2 &= 0. \end{aligned}$$

It follows from (28):

$$v(t, x) [1 - \beta_3 x_2 - \eta_4 x_1] = C_{11} x_1^2 + C_{12} x_1 x_2 - C_{22} x_2^2,$$

where

$$C_{11} = \eta_1 - \eta_4 K_{12}, \quad C_{12} = \gamma_1 - \beta_3 K_{12} - \eta_4 K_{22}, \quad C_{22} = \beta_3 K_{22}.$$

$$v(t, x) = \frac{C_{11}(t)x_1^2 + C_{12}(t)x_1x_2 - C_{22}(t)x_2^2}{1 - \beta_3x_2 - \eta_4x_1}, \quad t \in [0, T],$$

$$u(t, x) = -K_{12}(t)x_1 - K_{22}(t)x_2 + \frac{C_{11}(t)x_1^2 + C_{12}(t)x_1x_2 - C_{22}(t)x_2^2}{1 - \beta_3x_2 - \eta_4x_1}, \quad t \in [0, T]$$

provides stability in a finite time interval, i.e. $x(T) = 0$.

$$\begin{aligned} x_1(t) &= \frac{1}{(\mu_2 - \mu_1)\Delta_1} \{(\mu_2 e^{-\mu_1 t} - \mu_1 e^{-\mu_2 t})[(R_{11}R_{22}(0) - R_{12}R_{12}(0))x_{10} + \\ &+ (R_{12}R_{11}(0) - R_{11}R_{12}(0))x_{20}] + \varpi_n^2 (e^{-\mu_2 t} - e^{-\mu_1 t})[(R_{12}R_{22}(0) - R_{12}(0)R_{22})x_{10} + \\ &+ (R_{11}R_{22}(0) - R_{12}R_{12}(0))x_{20}]\}, \end{aligned}$$

$$\begin{aligned} x_2(t) &= \frac{1}{(\mu_2 - \mu_1)\Delta_1} \{(e^{-\mu_1 t} - e^{-\mu_2 t})[(R_{11}R_{22}(0) - R_{12}R_{12}(0))x_{10} + \\ &+ (R_{12}R_{11}(0) - R_{11}R_{12}(0))x_{20}] + (\mu_2 e^{-\mu_2 t} - \mu_1 e^{-\mu_1 t})[(R_{12}R_{22}(0) - R_{12}(0)R_{22})x_{10} + \\ &+ (R_{11}(0)R_{22} - R_{12}R_{12}(0))x_{20}]\}, \quad t \in [0, T]. \end{aligned}$$

$$C_{11} = \eta_1 - \eta_4 K_{12}, \quad C_{12} = \gamma_1 - \beta_3 K_{12} - \eta_4 K_{22}, \quad C_{22} = \beta_3 K_{22}.$$

$$v(t, x) = \frac{C_{11}(t)x_1^2 + C_{12}(t)x_1x_2 - C_{22}(t)x_2^2}{1 - \beta_3x_2 - \eta_4x_1}, \quad t \in [0, T],$$

$$u(t, x) = -K_{12}(t)x_1 - K_{22}(t)x_2 + \frac{C_{11}(t)x_1^2 + C_{12}(t)x_1x_2 - C_{22}(t)x_2^2}{1 - \beta_3x_2 - \eta_4x_1}, \quad t \in [0, T] \quad (29)$$

provides stability in a finite time interval, i.e. $x(T) = 0$.

The obtained control (29) at parametric excitation gives certain prescriptions for closing the current process and, although the mechanisms of the regulator may require detailed consideration, the proposed approach answers the question of providing the desired state [6]. From the latest results, it is not difficult to restore the initial designations of the variables of the respiratory chemostat [7].

In the article the stability of the respiratory chemostat on the finite time interval was studied, the task of stabilizing the biomedical system of the respiratory chemostat was solved, for cases of linear and quasilinear control systems and unsteady regime. The simulation of the process in a closed system for controlling the process of the respiratory chemostat for various values of the attenuation coefficient and its own angular frequency was carried out.

REFERENCES

- [1] Grodingz F. Teorija regulirovanija i biologicheskie sistemy. M.: Mir, 1966. 117 p.
- [2] La-Sall' Zh. Lefshec S. Issledovanie ustojchivosti prjamym metodom Ljapunova. M.: Mir, 1964. 37 p.
- [3] Abgarjan K.A. Vvedenie v teoriju ustojchivosti dvizhenija na konechnom intervale vremeni. M.: Nauka, 1992. 87 p.
- [4] Petrosjan P.L., Zaharov V.V. Matematicheskie modeli v jekologii. SPb.: Iz-vo SpbGU, 1997. 254 p.
- [5] Bijarov T.N., Kalimoldaev M.N., Kenesbaev S.M. Ustojchivost' i upravljaemost' nelinejnyh sistem // Dep. nauch. raboty. KazNIINKI. 1992. N 3911. K., 92, vyp. 2. 45 p.
- [6] Bellman R. Matematicheskie metody v medicine. M.: Mir, 1987. 200 p.
- [7] Wu M.-Y. On stability of linear time-varying system // Int. J. of systems Science. 1994. Vol. 15, N 2. P. 137-150.

Ж. К. Дюсембина

Л. Н. Гумилев атындағы Еуразия ұлттық университеті, Астана, Қазақстан

БИМЕДИЦИНАЛЫҚ ЖҮЙЕЛЕРДІ КВАЗИСЫЗЫҚТЫ БАСҚАРУ МОДИФИКАЦИЯСЫН ЗЕРТТЕУ

Аннотация. Тыныс алу хемостатының шектелген уақыт аралығындағы орнықтылығы зерттелген, экологиялық және биомедициналық жүйелерді зерттеудің сызықты және квазисызықты басқару жүйелері айқындалмаған түрде алынған.

Ағзадағы нормативтік процестерді зерттеудің осы түрі, өмір тіршілігін қамтамасыз ету үшін, техникалық жүйелерді жобалауда маңызды рөл атқарады. Және де бұл жүйе, оттегі мен көмірқышқыл газын, ағзадағы қысыммен реттеу ["тыныс алу хемостаты" жүйесіне] жүйесіне ғана байланысты емес, сонымен қатар басқа да өмірлік маңызды жүйелердің бірі-қатарын қамтиды. Бірінші кезекте бұл жүйеде "жүрек қантамырларының хемостатының" жүйесіне қатысты, ағзадағы таза қанның жылдамдығы арқылы, көміртегінің қысымын және оттегінің қалыпты жағдайға дейінгі деңгейіне жасаушы резервуарда жиналып, биологиялық тұрғыда өзін өзі реттейді.

Биологиялық өзін өзі реттеудің бұл күрделі гидродинамикалық жүйесі, көптеген параметрлері әр түрлі басқару сигналдарының үздіксіз басқаруының әсерінен, өз кезегінде, сыртқы ағзаның жағдайына тәуелді. Осы жүйенің реттеу режимі үздіксіз, күрт және кенеттен өзгеріп отыратын сыртқы әсерлері бар экстремалды сипатқа ие.

Түйін сөздер: тыныс алу хемостаты, басқару жүйесі, биологиялық жүйе, шектелген уақыт, орнықтылық,

Ж. К. Дюсембина

Евразийский национальный университет им. Л. Н. Гумилева, Астана, Казахстан

МОДИФИКАЦИЯ КВАЗИЛИНЕЙНОЙ СИСТЕМЫ УПРАВЛЕНИЯ БИМЕДИЦИНЫ

Аннотация. Исследована на устойчивость дыхательный хемостат на конечном интервале времени и решена задача стабилизации биомедицинской системы дыхательного хемостата, для линейной и квазилинейной систем управления. Изучение такого рода регуляционных процессов в организме играет важнейшую роль при разработке технических систем жизнеобеспечения, и это касается не только системы регулирования напряжения углекислого газа и кислорода (система "дыхательного хемостата"), но и ряда других жизненно важных систем. В первую очередь это относится к системе "сердечно-сосудистого хемостата", которая как биологическая система саморегулирования имеет задачей омывать тканевый резервуар свежей кровью с такой скоростью, чтобы поддерживать напряжение углекислого газа и кислорода в тканях на нормальном или близком к нему уровнях. Это сложная гидродинамическая система биологической саморегуляции, многочисленные параметры которой находятся под непрерывным воздействием различного рода управляющих сигналов, в свою очередь зависящих от внешних относительно организма условий и возмущений. Режим регуляции этой системы непрерывно подвергается резко и внезапно изменяющимся внешним воздействиям, имеющим экстремальный характер.

Ключевые слова: дыхательный хемостат, система управления, биологическая система, конечный интервал, устойчивость.

Сведения об авторе:

Дюсембина Жанар Какеновна – кандидат технических наук, Евразийский национальный университет им. Л. Н. Гумилева, Астана, Казахстан, zdyusembina@mail.ru

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://geolog-technical.kz/index.php/kz/>

Верстка Д. Н. Калкабековой

Подписано в печать 12.04.2017.

Формат 70x881/8. Бумага офсетная. Печать – ризограф.

16,7 п.л. Тираж 300. Заказ 2.