ҚАЗАҚСТАН РЕСПУБЛИКАСЫ ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

ХАБАРЛАРЫ

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

ГЕОЛОГИЯ ЖӘНЕ ТЕХНИКАЛЫҚ ҒЫЛЫМДАР СЕРИЯСЫ

СЕРИЯ ГЕОЛОГИИ И ТЕХНИЧЕСКИХ НАУК

SERIES
OF GEOLOGY AND TECHNICAL SCIENCES

4 (424)

ШІЛДЕ – ТАМЫЗ 2017 ж. ИЮЛЬ – АВГУСТ 2017 г. JULY – AUGUST 2017

ЖУРНАЛ 1940 ЖЫЛДАН ШЫҒА БАСТАҒАН ЖУРНАЛ ИЗДАЕТСЯ С 1940 г. THE JOURNAL WAS FOUNDED IN 1940.

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ ВЫХОДИТ 6 РАЗ В ГОД PUBLISHED 6 TIMES A YEAR

> АЛМАТЫ, ҚР ҰҒА АЛМАТЫ, НАН РК ALMATY, NAS RK

Бас редакторы

э. ғ. д., профессор, ҚР ҰҒА академигі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Редакция алкасы:

Абаканов Т.Д. проф. (Қазақстан)

Абишева З.С. проф., академик (Қазақстан)

Алиев Т. проф., академик (Әзірбайжан)

Бакиров А.Б. проф., (Қырғыстан)

Беспаев Х.А. проф. (Қазақстан)

Бишимбаев В.К. проф., академик (Қазақстан)

Буктуков Н.С. проф., академик (Қазақстан)

Булат А.Ф. проф., академик (Украина)

Ганиев И.Н. проф., академик (Тәжікстан)

Грэвис Р.М. проф. (АҚШ)

Ерғалиев Г.Х. проф., академик (Қазақстан)

Жуков Н.М. проф. (Казакстан)

Кенжалиев Б.К. проф. (Қазақстан)

Кожахметов С.М. проф., академик (Казахстан)

Конторович А.Э. проф., академик (Ресей)

Курскеев А.К. проф., академик (Қазақстан)

Курчавов А.М. проф., (Ресей)

Медеу А.Р. проф., академик (Қазақстан)

Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)

Нигматова С.А. проф. (Қазақстан)

Оздоев С.М. проф., академик (Қазақстан)

Постолатий В. проф., академик (Молдова)

Ракишев Б.Р. проф., академик (Қазақстан)

Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)

Сейтмуратова Э.Ю. проф., корр.-мушесі (Қазақстан)

Степанец В.Г. проф., (Германия)

Хамфери Дж.Д. проф. (АҚШ)

Штейнер М. проф. (Германия)

«КР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,

http://nauka-nanrk.kz/geology-technical.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: К. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

____ 2 ____

Главный редактор

д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Редакционная коллегия:

Абаканов Т.Д. проф. (Казахстан)

Абишева З.С. проф., академик (Казахстан)

Алиев Т. проф., академик (Азербайджан)

Бакиров А.Б. проф., (Кыргызстан)

Беспаев Х.А. проф. (Казахстан)

Бишимбаев В.К. проф., академик (Казахстан)

Буктуков Н.С. проф., академик (Казахстан)

Булат А.Ф. проф., академик (Украина)

Ганиев И.Н. проф., академик (Таджикистан)

Грэвис Р.М. проф. (США)

Ергалиев Г.Х. проф., академик (Казахстан)

Жуков Н.М. проф. (Казахстан)

Кенжалиев Б.К. проф. (Казахстан)

Кожахметов С.М. проф., академик (Казахстан)

Конторович А.Э. проф., академик (Россия)

Курскеев А.К. проф., академик (Казахстан)

Курчавов А.М. проф., (Россия)

Медеу А.Р. проф., академик (Казахстан)

Мухамеджанов М.А. проф., чл.-корр. (Казахстан)

Нигматова С.А. проф. (Казахстан)

Оздоев С.М. проф., академик (Казахстан)

Постолатий В. проф., академик (Молдова)

Ракишев Б.Р. проф., академик (Казахстан)

Сеитов Н.С. проф., чл.-корр. (Казахстан)

Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)

Степанец В.Г. проф., (Германия)

Хамфери Дж.Д. проф. (США)

Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

http://nauka-nanrk.kz/geology-technical.kz

© Национальная академия наук Республики Казахстан, 2017

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

____ 3 ____

Editor in chief

doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

Editorial board:

Abakanov T.D. prof. (Kazakhstan)

Abisheva Z.S. prof., academician (Kazakhstan)

Aliyev T. prof., academician (Azerbaijan)

Bakirov A.B. prof., (Kyrgyzstan)

Bespayev Kh.A. prof. (Kazakhstan)

Bishimbayev V.K. prof., academician (Kazakhstan)

Buktukov N.S. prof., academician (Kazakhstan)

Bulat A.F. prof., academician (Ukraine)

Ganiyev I.N. prof., academician (Tadjikistan)

Gravis R.M. prof. (USA)

Yergaliev G.Kh. prof., academician (Kazakhstan)

Zhukov N.M. prof. (Kazakhstan)

Kenzhaliyev B.K. prof. (Kazakhstan)

Kozhakhmetov S.M. prof., academician (Kazakhstan)

Kontorovich A.Ye. prof., academician (Russia)

Kurskeyev A.K. prof., academician (Kazakhstan)

Kurchavov A.M. prof., (Russia)

Medeu A.R. prof., academician (Kazakhstan)

Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)

Nigmatova S.A. prof. (Kazakhstan)

Ozdoyev S.M. prof., academician (Kazakhstan)

Postolatii V. prof., academician (Moldova)

Rakishev B.R. prof., academician (Kazakhstan)

Seitov N.S. prof., corr. member. (Kazakhstan)

Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)

Stepanets V.G. prof., (Germany)

Humphery G.D. prof. (USA)

Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

http://nauka-nanrk.kz/geology-technical.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Editorial address: Institute of Geological Sciences named after K.I. Satpayev

69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

____ 4 ____

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 4, Number 424 (2017), 133 – 141

UDC669.213.6

B. N. Surimbayev^{1, 2}, A. O. Baikonurova¹, L. S. Bolotova²

¹Kazakh National Research Technical University after K. I. Satpayev, Almaty, Kazakhstan, ²The Branch of the RSE «NC CPMS RK» State scientific-industrial association of industrial ecology "Kazmekhanobr", Almaty, Kazakhstan.

E-mail: surimbaev@gmail.com, a.baikonurova@yandex.kz, L bolotova@yahoo.com

PROSPECTS FOR THE DEVELOPMENT OF THE PROCESS OF INTENSIVE CYANIDATION OF GOLD-CONTAINING PRODUCTS IN THE REPUBLIC OF KAZAKHSTAN

Abstract. Data on the raw materials base of the gold mining industry of Kazakhstan, as well as the role of Kazakhstan in gold mining in the world and CIS countries are given. The history and development of the process of intensive cyanidation of gold-containing products in world practice is shown. The prospects of using this method in the Republic of Kazakhstan are shown. Technological schemes of intensive cyanidation are given depending on the material composition of the raw materials.

Key words: intensive cyanidation, gravity concentration, centrifugal concentrator, Acacia, ILRGekko, cyanidation, gold production.

In recent years, there has been a significant increase in the production of precious metals, particularly gold. This is due to the fact that gold is the main banking metal and is a currency reserve of many countries. This issue is especially topical because of the instability of the convertible world currencies, the fall in prices for securities and the consequences of the global financial crisis.

Gold in nature is found mainly in the native state, mainly in the form of small grains interspersed in quartz or contained in quartz sand. In small quantities, gold is found in sulphide ores of iron, lead and copper. Throughout history, mankind produced about 161 thousand tons of gold (estimate for 2011). If you fuse all this gold together, you get a cube with a side of about 20 m [1, 2].

Currently, 50% of gold is used in jewelry, 40% is used as an investment and the remaining 10% is used in industry [3].

The world gold production in 2015, according to Metals Focus, was 3211 tons [4]. The largest countries producing gold are listed in Table 1 and Figure 1.

Place	Country	Extraction, t	
		2014	2015
1	2	3	4
1	China	462,0	460,3
2	Australia	274,0	273,8
3	Russia	264,7	268,5
4	United States	210,0	214,0
5	Peru	171,1	170,5
6	South Africa	168,6	167,5
7	Canada	151,2	157,2

Table 1 – Twenty leading gold mining countries [4]

1	2	3	4
8	Mexico	112,7	133,2
9	Indonesia	93,8	113,0
10	Brazil	91,6	95,0
11	Ghana	106,3	94,7
12	Uzbekistan	83,5	85,5
13	Kazakhstan	49,2	63,7
14	Argentina	60,0	63,5
15	Papua New Guinea	60,7	58,4
16	Tanzania	50,8	51,7
17	Mali	52,8	50,1
18	Colombia	47,0	44,0
19	Philippines	40,4	41,1
20	Chile	44,5	41,1
	Others	558,1	564,4
	Global Total	3153	3211




Figure 1 – The largest gold producers in 2015

In 2014 Kazakhstan occupied the sixteenth place in the production of gold (49.2 tons), and in 2015, it took the thirteenth line, increasing gold production by 29% (63.7 tons) [4].

Table 2 shows gold production in the CIS countries in the periods from 2010 to 2015.

Extraction, t Country 2010 2011 2012 2013 2014 2015 Russia 203,1 211,6 229,3 248,5 264,7 268,5 Uzbekistan 71,0 70,6 80,0 81,0 83,5 85,5 63,7 Kazakhstan 30,3 36,8 40,0 42,4 49,2 Kyrgyzstan 19,0 19,7 11,4 20,2 19,3 18,7 Other 5,9 5,9 7,1 8,8 5,7 7,2 Total 329,0 344,7 366,6 399,2 423,9 445,3

Table 2 – Gold mining in the CIS countries [4]




Figure 2 – Graph of gold production in the CIS countries

As can be seen from Figure 2, Kazakhstan is the third largest gold producer among the CIS countries after Russia and Uzbekistan. And recently, the growth of gold production in Kazakhstan continues. The country's resources are estimated at 1.8 thousand tons, gold reserves are ~ 800 tons (according to 2009 data) [5].

Gold ore deposits have been identified in all regions of the republic, East (about 52.2%), North and Central Kazakhstan (30%) occupy the leading positions in terms of reserves, where 16 mining regions are located. The most important of them are: in the east - Kalbinsky and Rudno-Altaisky, in the north - Kokshetausky and Zhetigarinsky, in the south - Shu-Iliysky and Dzungarsky, in the west - Mugodzharsky, in the central part - Maikainsly and North Balkhash.

The raw material base of the gold mining industry in Kazakhstan is represented mainly by small (with reserves of up to 25 tons) and medium (from 25 to 100 tons) deposits. According to some sources, there are 122 indigenous gold ore deposits, 81 integrated and 34 placer mines [6, 7].

Gold ore deposits in Kazakhstan are characterized by low metal content in the ore, as well as the presence of a significant proportion of refractory ore. At the same time, the comparatively favorable mining technical conditions of extraction make it possible to keep its production costs at the level of the world average.

Easy enriched and oxidized ores, which are the main resource of gold mining companies, are already running low and half of all gold reserves are concentrated in the 8 largest complex fields –Bakyrchik, Vasilkovskoye, Mizek, Suzdalskoe, Bolshevik, Akbakay, Bestobe and Zholymbet. In them, gold-bearing ores are generally classified as stubborn [6].

Currently, licenses are issued for about half of the number of deposits, of which about 30 are in service. Some industrial enterprises extract gold as a by-product in the processing of polymetallic ores. The "KAZ Minerals" company (formerly Kazakhmys), the leading copper producer in the country, also simultaneously extracts gold from the ore. In 2012, gold production amounted to 3.6 tons (with a content of 0.66 g/t), in 2013 - 3.2 tons (at a content of 0.61 g/t) [8, 9].

The largest gold mining facility in Kazakhstan is the Vasilkovskoye deposit, located 17 km northwest of Kokshetau. The proven reserves of deposits amount to 370 tons of gold, with an average ore grade of 2.8 g/t. The deposit has been under development since 1979. Since 1991, at the Vasilkovskoe Ore Mining and Processing Enterprise, the processing of oxidized ore was carried out by heap leaching. The capacity of the heap leach plant was 1 million tons of ore per year with an average annual production of 600 to 900 kg of gold. In 2007, the deposit began construction of a large gold recovery plant with modern ore processing technology. At full design capacity, the mine should produce 15 tons of gold per year. Capital investments amounted to about 700 million dollars. The factory was built in two years. In 2010, at the suggestion of the Head of state VasilkovskyGOK was renamed into the "AltyntauKokshetau" company [8, 10].

The second place is occupied by the Bakyrchik deposit located in the northeast of Kazakhstan (about 100 km from Semipalatinsk) [11]. In the second largest gold deposit of Kazakhstan, gold mining is still not conducted due to the high persistence of the ore, high content of arsenic and organic carbon in the ore. Plans for processing such ore by bioleaching have not been implemented, since this technology requires high costs [7, 12, 28].

The following in order of industrial importance can be noted the Aksu, Bestobe, Zholymbet. All of them are worked underground. The ores in the deposits are represented as quartz veined, and in the form of mineralized zones. The mines are located within a radius of 100 km from the central base, located in Stepnogorsk (Akmola region). The richest and easily accessible ores are worked out, in connection with this, in the period of low gold prices there was a decline in production. However, the significant gold reserves of these deposits are more than 100 tons, and the predicted reserves are 400 tons [8, 28].

Despite the fact that Kazakhstan's major and medium size fields are currently mostly exploited, the country has a significant number of small deposits that require additional exploration (Table 3).

Location	Gold Deposits	
North Kazakhstan	Vasilkovskoye, Varvarinskoye, Uzboi, Symbat, Komarovskoye, Elevatornoye, Akkarginskoye, Zhetygorinskoye	
Central Kazakhstan	Aksu, Zholymbet, Bestobe, Maikain, Kvartsytovye Gorki, Ushoky, Yenbekshi, Pustynnoye	
East Kazakhstan	Bakyrchik, Suzdalskoye, Sekosovskoye, Bolshevik, Vasiliyevskoye, Ridder-Sokolnoye, Zhanan, Akzhal, Kaskabulak	
South Kazakhstan	Akbakai, Altyntas, Dalabai, Aksakal-Beskempir, Mynaral, Zharkulak, Karamurun, Arkharly, Kumysty	
West Kazakhstan	Yubelienoye	

Table 3 – Main gold deposits in Kazakhstan [6]

Therefore, the development of technology for the effective enrichment of gold-containing raw materials and the development of research in this direction is timely and important.

Methods of ore dressing consist of the ore pretreatment operation (crushing, screening, sorting, grinding) and methods of beneficiation of prepared ore (gravity, flotation, magnetic separation, etc.). Metallurgical methods for processing ores and concentrates include cyanidation (or other leaching techniques), amalgamation, roasting, sorption methods for extracting gold from pulps and solutions, extractive methods, hydrochlorination, autoclave oxidation [13].

It is known that the most common process of extracting gold from poor ores is cyanidation. This process is based on the selective leaching of gold or other noble metal by aqueous solutions of alkaline cyanides: sodium, potassium or calcium. The resulting solution containing dissolved gold is sent for processing by various methods to obtain a high-quality commodity product in the form of Dore gold in ingots. The resulting alloy is sent to the refinery to produce gold of the required purity [14].

In the practice of gold production for the extraction of large grains of free metal, gravity enrichment is used, which is the oldest method of processing mineral raw materials [15]. For a long period of time this method has undergone changes - from simple washing and separation of grains on the inclined plane to the use of centrifugal concentrators. The method of separation of mineral grains by density in gravity devices is simple and can enrich placer gold in remote areas without the existing infrastructure. In addition, this method does not require the use of chemical reagents and is characterized by low energy intensity [16, 17]. The need to allocate large gold, which is present in almost all ores, is determined by the following reasons [18-20]:

- the duration of dissolution of large particles during cyanidation is increased;
- large particles of gold will accumulate in grinding and classifying equipment;
- particles of large gold in the grinding cycle (mill-classifier) are overwritten.

To prevent the accumulation of gold in the classifier and overwriting in the mill, free gold is continuously withdrawn from the circulating load by the gravitational enrichment method.

New technological schemes for the enrichment of gold-containing products, which include gravity on centrifugal concentrators, allow to obtain concentrates rich in gold content (>100 g/t) with a relatively small yield.

Hydrometallurgical processing of rough gravity concentrates is carried out by cyanidation. The essence of the method lies in the leaching (dissolution) of gold using solutions of cyanide salts in the presence of an oxidizer (oxygen). Factors that affect the cyanidation process can be conditionally divided into two groups: depending on the material composition (gold size, gold form, gold surface and presence of impurities) and technological ones that can be controlled - mixing, temperature, pressure, pH value, viscosity of pulps [21].

Hydrometallurgical processing of these concentrates by traditional methods of cyanidation is difficult due to the lack of equipment of low productivity and significant losses of metal with a cake.

Until recently, the only acceptable method of processing them was considered deep debugging (cleaning), followed by melting the resulting "gold heads" on metal ingots. However, special devices have now been created that permit the leaching of large grains of metallic gold by cyanide solutions.

Since the early 1990s, the introduction of centrifugal concentrators (Knelson, Falcon, etc.) in many plants has significantly increased the extraction of gold into gravity concentrates (by 1-3%), including the processing of ores with a high sulfide content [22-25]. However, standard methods for processing gravity concentrates allow gold recovery at a level of no more than 70% [26]. To solve this problem, intensive cyanidation plants have been tested, which make it possible to achieve a high extraction of gold from gravity concentrates for an acceptable time for the general technological cycle (6-24 h).

In the process of intensive cyanidation, solutions with a higher concentration of cyanide, protective alkali with the addition of soluble compounds of lead (II) and an oxidizer are used.

For the first time in the world, the intensive cyanidation method for the processing of gravity concentrates was applied at the South Africa plant (Western-Holdings) in 1977 and then at the gold-processing plant in East-Gold in 1980. As leach reactors, column type plants with mechanical mixing, equipped with a circulator or revolving arms, were used [27, 28].

But, until recently, interest in the intensive cyanidation process has faded, due to a number of reasons. Firstly, the largest corporations did not consider the gravitational enrichment as a strategic direction for the development of technology for processing South African gold-containing ores. Secondly, the use of agitators with mechanical mixing caused technical problems associated with the rapid deterioration of equipment. Thirdly, there were no opportunities to automate the process, which created the prerequisites for the organization of gold theft [22].

The renewed interest in the process of intensive cyanidation occurred relatively recently in connection with the use of reactors and new type installations. Thus, the Acacia intensive cyanidation reactor was developed by AngloGoldAustralia for the Union-Reef plant (South Africa) in 1999 [25]. In mid-2003, these reactors were used in 27 plants of leading foreign gold mining companies: in Australia – 11; in Zimbabwe – 5; in Canada – 3; inSouth Africa, Papua New Guinea and Mongolia –by 1; in Russia – 5 [29].

Acacia is a complex compact installation, the core of which is a cone-shaped reactor with a suspended layer of concentrate due to the upward flow of the solution. Due to the high concentration of reagents in the solution (2-3% NaCN, 0.2-0.35% NaOH, 0.1-1.0% Leach - accelerator), intensive cyanidation leads to the dissolution of gold both free and rising into the crystal lattice of sulfide minerals. This makes it possible to reduce the number of pieces of equipment in the subsequent processing stages, to reduce the gold content in the tailings, and also to increase the overall gold recovery by 1.0-1.5% [30, 31].

Leaching of gold by alkaline cyanide solutions in the presence of oxygen or air is a widely used process for extracting gold from mineral raw materials. The gold is oxidized and dissolved in the presence of cyanide to form the complex ion $[Au(CN)_2]$ according to the Bodlander (1) and (2) and Elsner (3) equations, the latter is the sum of equations (1) and (2)[32-35]:

$$2Au + 4CN^{-} + O_{2} + 2H_{2}O \rightarrow 2Au(CN)_{2}^{-} + 2OH^{-} + H_{2}O_{2},$$
 (1)

$$2Au + 4CN^{-} + H_{2}O_{2} \rightarrow 2Au(CN)_{2}^{-} + 2OH^{-},$$
 (2)

$$4Au + 8CN^{-} + O_2 + 2H_2O \rightarrow 4Au(CN)_2^{-} + 4OH^{-}.$$
 (3)

The behavior of gold during cyanidation depends on a number of factors, of which the main is the relationship of gold to ore and rock-forming minerals. Gold can be as in a free state, and in the form of splices with minerals. Any of these gold conditions can affect its under-extraction during dissolution and cause an increased consumption of reagents [36].

Excessive oxidation of sulphides during gold leaching can also lead to gold passivation due to the formation of gold surface coatings by oxidation products [35].

The presence of sulfides such as marcasite, pyrrhotite or chalcocite in cyanate pulp often inhibits the dissolution of gold, forming a protective film on the surface of gold [21].

To intensify the process of gold leaching, chemical additives are used. Chemical additives, depending on their nature, can be divided into inorganic and organic. Cyanidation accelerators under the trade names LeachAid, LeachWell and others are a mixture of organic and inorganic sodium salts (93-99%), lead nitrate (1-5%) and water (up to 5%). Made in the form of small granules of slightly yellow color, odorless LeachWell or with a weak smell of caramel LeachAid are stable up to a temperature of 300 °C. The density of these reagents is 1 g/cm³, the solubility in water is 35-40 g/dm³[21, 31].

Under the conditions of standard cyanidation, the presence in the solution of the reagent-accelerator makes it possible to increase the dissolution rate of gold and the degree of its extraction from the gravitational concentrate and to shorten the leaching time [31].

In 1997, in Australia (GekkoSystems), an intensive cyanidation unit was developed – InLineLeachReactor (ILR), originally introduced at a number of African factories of AshantiGoldfields, and is currently used in more than 20 foreign factories [31, 37].

The ILR reactor is also a simple apparatus for intensive cyanidation of gravity concentrates. The ILR reacts according to the semi-submerged layer principle and represents a horizontally mounted drum rotating at a low speed, with a set of special partition walls and an aeration system for maximum leaching. In addition to the proven reactors for intensive cyanidation of Acacia and ILRGekko, there are other devices of different design and characteristics.

Many domestic gold mining companies, still using traditional technology of gold enrichment, begin to show interest in the process of intensive cyanidation. Thus, the Acacia (Knelson/ConSep) intensive cyanidation reactors are installed and operate at the fields Pustynnoe and Akbakay for the enrichment of gold-bearing concentrates [38].

Intensive cyanidation, depending on the material composition, can be used in a variety of technological schemes. Figure 3 shows a flow chart of intensive cyanidation for the processing of easily cyanidized raw materials. According to the presented technology, concentrates after gravity concentration are sent to the intensive cyanidation process, in which, productive solutions are processed by electrolysis methods. Thecakeissenttothetailingdump.




Figure 3 - Technological scheme of intensive cyanidation for the processing of easily cyanidized raw materials

For oxidized ores with a small content of sulphides, a process scheme is used in which the cake after intensive cyanidation is returned to the process for complete gold recovery (Figure 4).




Figure 4 – Technological scheme of intensive cyanidation for oxidized ores

In the scheme for processing sulphide ores, the cake after intensive cyanidation is directed to sorption leaching (Figure 5).




Figure 5 – Technological scheme of intensive cyanidation for sulphide ores

The use of the intensive cyanidation process allows to increase the processing of gold-containing ores by introducing more branched technological schemes.

The data presented in the review show that intensive cyanidation represents a prospect for the processing of gold-containing materials, because it shortens the duration of the cyanidation process and reduces the loss of gold, thereby achieving a high recovery of gold from the gravity concentrate into the solution.

REFERENCES

- [1] Surimbayev B.N., Bolotova L.S., Baikonurova A.O., Mishra B. Study of process cyanide leach gold from ore and gravity concentration tailings sulfide and oxide ores [Izuchenie processa cianidnogo vyshhelachivanija zolota iz rudyihvostov gravitacionnogo obogashhenija sul'fidnyhiokislennyhrud]. Bulletin of National Academy of Sciences of the Republic of Kazakhstan.Vol. 4. Number 362 (2016). P. 260-266. ISSN: 1991-3494 (In Russian)
 - [2] The cost and reserves of gold in the world. Analytics.http://ria.ru/infografika/20110824/422749631.html
- [3] Soos, Andy. Gold Mining Boom Increasing Mercury Pollution Risk. http://oilprice.com/Metals/Gold/Gold-Mining-Boom-Increasing-Mercury-Pollution-Risk.html
- [4] Metals Focus Gold Focus 2016. London. 2016. P. 15-19. ISBN: 978-0-9935876-0-3. http://info.sharpspixley.com/uploads/GoldFocus2016.pdf
- [5] Donskih A. Barrier for gold: What prevents the noble metal from becoming the leading financial market. The newspaper is Kazakhstanskaya Pravda. 2009. 25 June. P.8 (In Russian)
- [6] Michael E. Wilson, Elena Lee. Kazakhstan's Gold Mining Sector and the New Regulation on Gold Sales. Alchemistissuesixtysix. P. 12-15(in Eng.)
- [7] BulenbaevM.Zh. Development of technology for extracting gold from persistent gold-bearing raw materials [Razrabotka tehnologi i izvlechenijazolotaizupornogozolotosoderzhashhegosyr'ja]. Dissertation.Almaty. Kazakhstan. 2015. P. 13-14 (In Russian)
- [8] Verhozin S.S. Gold Mining Industry of Kazakhstan [Zolotodobyvajushhajapromyshlennost' Kazahstana]. https://zolotodb.ru/news/11194
 - [9] Kazakhmys PLC.Annual Report and Financial Statements for 2013.http://www.kazminerals.com/pdf/ara_2013_ru.pdf
 - [10] Altyntau Kokshetau.http://www.altyntau.com/
 - [11] Gold of Kazakhstan: a brief review.http://www.investkz.com/journals/38/236.html
- [12] Speech Madiev B.M. at a seminar-meeting in the Ministry of Industry and New Technologies of the Republic of Kazakhstan. The state of the raw material base of the gold mining industry, the direction and ways of development of geological exploration for gold in the Republic of Kazakhstan.04.02.2014. http://www.parasat.com.kz/index.php?id=899&L=2
- [13] Bajysbekov Sh. Combined cyanide-free processing technology of resistant gold-bearing weathering crusts [Kombini-rovannajabescianidnajatehnologijapererabotkiupornyhzolotosoderzhashhihrudkoryvyvetrivanija]. Dissertation. Almaty. Kazakhstan. 2008. P. 260(In Russian)
- [14]MarsdenJ., HouseI. TheChemistryofGoldExtraction.SocietyforMining, MetallurgyandExploration. USA. 2006. P. 48-50. ISBN-13: 978-0-87335-240-6 / ISBN-10: 0-87335-240-8 (in Eng.)
- [15]Meretukov M.A. Gold: the origin of mining, metallurgy and technology [Zoloto: zarozhdeniegornogodela, metallurgii i tehnologii]. Moscow. RudaiMetally. 2008. P. 180. ISBN: 978-5-98191-042-5(In Russian)
- [16] Verhoturov M.V. Gravityconcentrationmethods [Gravitacionnyemetodyobogashhenija]. Moscow. 2006. P.7-18. ISBN: 5-317-01710-6 (In Russian)
- [17]Surimbayev B.N., Baikonurova A.O., Bolotova L.S. Investigation of the process of gravity concentration of gold-containing sulfide ores. [Issledovanieprocessagravitacionnogoobogashhenijazolotosoderzhashhihsul'fidnyhrud].Reports of the National Academy of Sciences of the Republic of Kazakhstan. Vol. 3. N 313 (2017). P. 55-60. ISSN: 2224-5227 (In Russian)
- [18] Chryssoulis, S.L. and McMullen, J. Mineralogical Investigation of Gold Ores. In: Adams, M.D., Ed., Advances in Gold Ore Processing. Elsevier.The Netherlands.2005. P. 21-67. DOI: 10.1016/S0167-4528(05)15002-9(in Eng.)
- [19] Meretukov M.A.Gold. Chemistry, Mineralogy, Metallurgy[Zoloto.Himija, mineralogija, metallurgija].Moscow. RudaiMetally. 2008. P. 520. ISBN: 978-5-98191-043-2(In Russian)
- [20] Lodejshhikova V.V. Technology and technology of extracting gold from ores abroad [Tehnika i tehnologija izvlechenija zolotaizrudzarubezhom]. Moscow. Metallurgy. 1973. P. 288(In Russian)
- [21] Evdokimov A.V. Investigation of the intensive cyanidation process of gold-containing gravity concentrates [Issledovanie processaintensivnogocianirovanijazolotosoderzhashhihgravitacionnyhkoncentratov]. Dissertation. Irkutsk. Russia. 2012. P. 9-20 (InRussian)
- [22] Zaharov B.A., Meretukov M.A.Gold: resistant ores[Zoloto: upornyerudy]. Moscow. RudaiMetally. 2008. P.296-300. ISBN: 978-5-98191-068-5(In Russian)
- [23] Ancia, P. H., Frenay, J., Dandois, P. H., 1997. Comparison of the Knelson and Falcon centrifugal separators. In: Richard M. Mozley International Symposium (in Eng.)
- [24] Longley R., McCallum A., Katsikaris N. Intensive cyanidation: onsite application of the InLine Leach Reactor to gravity gold concentrates. Minerals Engineering. 2003. V. 16. P. 411-419. DOI:10.1016/S0892-6875(03)00054-2(in Eng.)
- [25] Campbell J. & Watson B. Gravity Leaching with the ConSep ACACIA Results from AngloGold Union Reefs. Eigth Mill Operators Conference, 21-23 July. 2003.(in Eng.)
- [26] Laplante A., Staunton W. // Pros. 5th Int. Symp.: Hydrometallurgy 2003 / Vancouver, Canada. 2003. August 24-27. V. 1. P. 65-74 (in Eng.)
- [27]Davidson, J. D., Brown, G. A., Schmidt, C. G., Hanf, N. W., Duncanson, D., Taylor, J. D. The intensive cyanidation of gold-plant gravity concentrates. Journal of the SouthAfricanInstitute ofMining andMetallurgy.Vol. 78. 1978. P. 146-165 (in Eng.)
- [28] Dewhirst R.F., Moult S.P., Coetzee J.A. Intensive cyanidation for the recovery of coarse gold. Journal of the SouthAfricanInstitute of Mining and Metallurgy. Vol. 84. 1984. P. 159-163 (in Eng.)
- [29] Car'kov V.A. Experience of gold extracting enterprises in the world[Opytrabotyzolotoizvlekatel'nyhpredprijatijmira]. Moscow. RudaiMetally.2004. P. 18-75. ISBN: 5-98191-006-2(In Russian)
 - [30] Gravity leaching the Acacia reactor. http://knelsongravity.xplorex.com/sites/knelsongravity/files/reports/report36s.pdf

- [31] Meretukov M.A. Development of intensive cyanidation of gold-containing gravity concentrates [Razvitie processa intensivnogo cianirovanijazolotosoderzhashhihgravitacionnyhkoncentratov]. Non-ferrous metals.2005. №2. P. 39-42. ISSN: 0372-2929.(In Russian)
- [32] Ling P., Papangelakis V.G., Argyropoulos S.A. et al. An Improved Rate Equation for Cyanidation of a Gold Ore // Canadian Metallurgical Quarterly, 1996. Vol. 35, N. 3, P. 7 (in Eng.)
- [33]Kondos P.D., Deschenes G., Morrison R.M. Process Optimization Studies in Gold Cyanidation // Hydrometallurgy. 1995. V. 39. P. 235-250 (in Eng.)
- [34] de Andrade Lima, L.R.P., Hodouin D. A Lumped Kinetic Model for Gold Ore Cyanidation // Hydrometallurg. 2005. V. 79 (in Eng.)
- [35] Turysbekova G.S., Meretukov M.A., Bektaj E.K. Gold: Innovations in Chemistry and Metallurgy [Zoloto: innovacii v himiiimetallurgii]. Almaty. 2015. P. 117. ISBN:978-601-228-838-4 (In Russian)
 - [36]Karimi P. et al. // Int. J.Miner.Proc. 2010. V. 95. P. 68-77(in Eng.)
 - [37]InLine Leach Reactor.http://www.gekkos.com/equipment/inline-leach-reactor
 - [38] AltynalmasAKJSC. Investment memorandum.Almaty. 2015. http://www.kase.kz/files/emitters/ALMS/almsf7_2015.pdf

Б. Н. Суримбаев^{1, 2}, А. О. Байконурова¹, Л. С. Болотова²

¹Казахский национальный исследовательский технический университет им. К. И. Сатпаева, Алматы, Казахстан, ²Филиал РГП «НЦ КПМС РК» Государственное научно-производственное объединение промышленной экологии «Казмеханобр», Алматы, Казахстан

ПЕРСПЕКТИВЫ РАЗВИТИЯ ПРОЦЕССА ИНТЕНСИВНОГО ЦИАНИРОВАНИЯ ЗОЛОТОСОДЕРЖАЩИХ ПРОДУКТОВ В РЕСПУБЛИКЕ КАЗАХСТАН

Аннотация. Приведены данные по сырьевой базе золотодобывающей промышленности Казахстана, а также роль Казахстана по добыче золота в мире и странах СНГ. Показана история и развитие процесса интенсивного цианирования золотосодержащих продуктов в мировой практике. Показана перспективность использования данного метода в Республике Казахстан. Даны технологические схемы интенсивного цианирования в зависимости от вещественного состава сырья.

Ключевые слова: интенсивное цианирование, гравитационное обогащение, центробежный концентратор, Акация, Гекко, цианирование, добыча золота.

Б. Н. Сүрімбаев^{1, 2}, Ә. Ө. Байқоңырова¹, Л. С. Болотова²

¹Қ. И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті, Алматы, Қазақстан, ²«ҚР МШКҚӨҰО» РМК Мемлекеттік өнеркәсіптік экология ғылыми-өндірістік бірлестігі филиалы «Қазмеханобр», Алматы, Қазақстан

ҚАЗАҚСТАН РЕСПУБЛИКАСЫНДАҒЫ АЛТЫНҚҰРАМДЫ ӨНІМДЕРДІ ҚАРҚЫНДЫ ЦИАНИРЛЕУ ҮРДІСІНІҢ ДАМУЫНЫҢ БОЛАШАҒЫ

Аннотация. Мақалада Қазақстанда алтын өндіру өнеркәсіп шикізат негізінің көрсеткіштері, сонымен қатар әлемдегі және ТМД елдері арасындағы алтын өндірісі бойынша Қазақстанның рөлі көрсетілген. Әлемдік тәжірибедегі алтынқұрамды өнімдердің қарқынды цианирлеу процесінің тарихы және дамуы көрсетілген. Аталған әдістің Қазақстан Республикасындағы қолдану болашағы көрсетілген. Шикізаттың заттай құрамына тәуелді қарқынды цианирлеу үрдісінің технологиялық схемалары сипатталған.

Түйін сөздер: қарқынды цианирлеу, гравитациялық байыту, ортадан тепкіш концентратор, Акация, Гекко, цианирлеу, алтын өндірісі.

Information about authors:

Surimbayev Bauyrzhan Nurzhanovich – PhD student, Kazakh National Research Technical University after K . I. Satpayev, Almaty, Kazakhstan; Research scientist, laboratory of noble metals, The Branch of the RSE «NC CPMS RK» State scientific-industrial association of industrial ecology "Kazmekhanobr", Almaty, Kazakhstan, surimbaev@gmail.com

Baikonurova Aliya Omirkhanovna – Doctor of Technical Sciences, Professor, Kazakh National Research Technical University after K. I. Satpayev, Almaty, Kazakhstan, a.baikonurova@yandex.kz

Bolotova Lyudmila Sergeyevna – Candidate of Chemical Sciences, Head of the laboratory of noble metals, The Branch of the RSE «NC CPMS RK» State scientific-industrial association of industrial ecology "Kazmekhanobr", Almaty, Kazakhstan, L_bolotova@yahoo.com

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see http://www.elsevier.com/publishingethics and http://www.elsevier.com/journal-authors/ethics.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see http://www.elsevier.com/postingpolicy), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service http://www.elsevier.com/editors/plagdetect.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

http://geolog-technical.kz/index.php/kz/

Верстка Д. Н. Калкабековой

Подписано в печать 11.07.2017. Формат 70x881/8. Бумага офсетная. Печать – ризограф. 18,2 п.л. Тираж 300. Заказ 4.