

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный исследовательский
технический университет им. К. И. Сатпаева

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Kazakh national research technical university
named after K. I. Satpayev

ГЕОЛОГИЯ ЖӘНЕ ТЕХНИКАЛЫҚ ҒЫЛЫМДАР СЕРИЯСЫ



СЕРИЯ ГЕОЛОГИИ И ТЕХНИЧЕСКИХ НАУК



SERIES OF GEOLOGY AND TECHNICAL SCIENCES

1 (427)

ҚАҢТАР – АҚПАН 2018 ж.
ЯНВАРЬ – ФЕВРАЛЬ 2018 г.
JANUARY – FEBRUARY 2018

ЖУРНАЛ 1940 ЖЫЛДАН ШЫҒА БАСТАҒАН
ЖУРНАЛ ИЗДАЕТСЯ С 1940 г.
THE JOURNAL WAS FOUNDED IN 1940.

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Б а с р е д а к т о р ы

э. ғ. д., профессор, ҚР ҰҒА академигі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Р е д а к ц и я а л қ а с ы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., академик (Қазақстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қырғыстан)
Беспәев Х.А. проф. (Қазақстан)
Бишимбаев В.К. проф., академик (Қазақстан)
Буктуков Н.С. проф., академик (Қазақстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Ерғалиев Г.К. проф., академик (Қазақстан)
Жуков Н.М. проф. (Қазақстан)
Кенжалиев Б.К. проф. (Қазақстан)
Қожахметов С.М. проф., академик (Қазақстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., академик (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Нигматова С.А. проф. (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Қазақстан)
Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)
Сейтмуратова Э.Ю. проф., корр.-мүшесі (Қазақстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«ҚР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2018

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н н а я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., академик (Казахстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Беспаяев Х.А. проф. (Казахстан)
Бишимбаев В.К. проф., академик (Казахстан)
Буктуков Н.С. проф., академик (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Ергалиев Г.К. проф., академик (Казахстан)
Жуков Н.М. проф. (Казахстан)
Кенжалиев Б.К. проф. (Казахстан)
Кожаметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., академик (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Нигматова С.А. проф. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Казахстан)
Сейтов Н.С. проф., чл.-корр. (Казахстан)
Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© Национальная академия наук Республики Казахстан, 2018

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

E d i t o r i a l b o a r d:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., academician (Kazakhstan)
Agabekov V.Ye. academician (Belarus)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Bespayev Kh.A. prof. (Kazakhstan)
Bishimbayev V.K. prof., academician (Kazakhstan)
Buktukov N.S. prof., academician (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadjikistan)
Gravis R.M. prof. (USA)
Yergaliev G.K. prof., academician (Kazakhstan)
Zhukov N.M. prof. (Kazakhstan)
Kenzhaliyev B.K. prof. (Kazakhstan)
Kozhakhmetov S.M. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., academician (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Nigmatova S.A. prof. (Kazakhstan)
Ozdoev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Rakishev B.R. prof., academician (Kazakhstan)
Seitov N.S. prof., corr. member. (Kazakhstan)
Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 1, Number 427 (2018), 99 – 106

UDC 910.4:633.32 (57.047/58.009)

**T. K. Salikhov¹, M. K. Tynykulov¹, I. F. Kostikov²,
Zh. Sh. Urgaliev¹, G. M. Salhozhaeva¹, Sh. E. Arystanova¹**

¹L. N. Gumilyov Eurasian National University, Astana, Kazakhstan,

²Sh. Ualikhanov Kokshetau State University, Kokshetau, Kazakhstan.

E-mail: salikhov_tk@enu.kz

THE BIOLOGICAL CHARACTERISTICS OF GRAIN PRODUCTION AND SILAGE FROM CORN (ZEA MAYS)

Abstract. Improvement of technology of cultivation of maize in the Northern Kazakhstan was based on the principles of soil-protective arable farming and harvesting silage was the main purpose. The advanced technology with the use of early-maturing hybrids (FAO (Food and Agriculture Organization) 150-199) on the basis of mini-mizing of the technological cycle has been developed in the North Kazakhstan Research Institute of Agriculture in the period from 1987 to 1997. The average output of absolutely dry substance for this period was 4.8 t/ha, that is 45.1% higher than in the previous period (1977–1986). However, the advanced technology based on the cultivation of the only silage crop (monoculture) cannot exclude a sharp decrease of the yields in droughty years and in years when the summer is short and cool. The silage conveyor as an alternative to monoculture of maize was substantiated by the selection of crops with different biological requirements for growth conditions. The yields of the silage conveyor crops (sunflower-maize-sweet sorghum) were 5.14 t/ha (absolutely dry substance), that is 0.73 t/ha higher than the yields of monoculture of maize in the period from 2007 to 2009.

Key words: maize, silage, fodder, grain, early maturing hybrids, dry substance, the cultivation technology, methods of cultivation, crop productivity, silo conveyor, monoculture, the Northern Kazakhstan.

Introduction. Maize cultivation in the northern regions of Kazakhstan originates from the 1950s when the maize began to sow to produce silage. The large areas were sowed for the first time (more than 75 thousands of hectares) in 1954 and the areas about 1.5 mln hectares, which is 2/3 of the total area of Kazakhstan, were sowed in the next 5 years [1, 2].

Large-scale introduction of maize and implementation of tilled farming system were carried out simultaneously. The maize field was the main one in crop rotations in this farming system [3].

The introduction of maize was carried out rapidly and over the large areas, so the zonal features of the new region of cultivation were not taken into account. Agricultural techniques, the system of agricultural implements and hybrids of maize were carried over unchanged. Agro techniques were based on late-ripening hybrids and the use of moldboard plowing, disk plough-harrows and checkrow planting ensured the ability of two inter-row processing in two directions during the period of vegetation.

Such unreasonable transfer of a tilled technology into the steppe zone, not protected from destructive influence of wind, led to the widespread deflation and systematic damage of the plants in the fields [4, 5].

The agricultural science faced a challenge of development of soil-protecting activities to overcome destructive influence of drought, wind erosion. These methods would guarantee the preservation of soil fertility. However, there were positive results. A practical expediency and economic efficiency of maize cultivation to harvest silage on the basis of imported seeds was proven during the period of introduction of maize in the Northern Kazakhstan [6-8].

Dairy farming was now based on a fundamentally new type of feeding in virgin areas. The basis of the ration was silage and concentrates [9]. Scientific research of that period lasted until 1986. Soil-

protective technologies were specified for each agricultural zone. The main tasks of the scientific research were to protect the plants in the fields from wind erosion and a decrease in the amplitude of fluctuations of the yields of maize. Researchers of the Northern Kazakhstan have proven that maize is not a tilled crop and there is no need for continuous loosening of row-spacing [10, 11].

Agro-physical parameters of the soil after plowing do not meet the biological requirements for the growth of maize. This fact was experimentally determined for the first time in the conditions of Northern Kazakhstan. When the density of the soil in the rooting zone is 1.05-1.1 g/cm³, the optimum water, air and food mode for the root system is created [12, 13], whereas after the plowing it is 0.9-0.95 g/cm³. Full substantiation of this theory was given by scientists of Agrophysical Scientific and Research Institute (Russian Federation) and in the countries of Western Europe [14, 15].

Materials and methods. The trends in change of the yields for a period of 1969-1986 were determined on the basis of statistical reporting in the administrative districts, which are included in the appropriate agricultural zone. Mathematical processing of data was performed by the method of equal squares [16].

The analysis of the chemical composition of the biomass was performed according to the conventional technique:

- the protein content - by the method of Kjeldahl;
- the determination of cellulose - by the method of Ganneberg-Shtoman;
- the determination of sugars - by the centrifuge micromethod of Bertrand-Bieri;
- the determination of crude fat - by extracting with ethyl ether;
- the definition of carotene - on the SF-596 spectrophotometer.

Results and discussion. Soil-protective farming system included new crop rotations with the "corn-wheat" crop group. The highest yield (absolutely dry substance) per 1 ha was obtained in this group.

Wide-row method of sowing (70 cm) has become an alternative to a checkrow sowing, where the integrated protection of maize was carried out by applying of soil herbicides and surface treatment the most littered plots on the field. A distinctive feature of the soil-protective technologies that had already been developed was at the zonal level, to minimize a technological cycle by consolidating of manufacturing operations, increasing the width of capture of units and decreasing of the depth of processing.

We carried out a comparative assessment of energy consumption for a tilled technology (during the period OF 1969-1976) and for a soil-protective technology (during the period OF 1977-1986) at the experimental field of the North-Kazakhstan Scientific and Research Institute of Agriculture. The experimental field is located in agricultural steppe zone of the Northern Kazakhstan. Abandoning the plough began to use the cultivator and this led to increased productivity of labor and to reducing of material resources. The maize seeder has been upgraded for stable operation after treatment of the ground with a cultivator and performed a number of operations in a single pass: seed placement, fertilizing, introduction of soil herbicides and soil loosening in the row zone [17].

Thus, the number of passes on the field has decreased almost by half (table 1).

A labor costs per unit of area has decreased respectively. The average yield for the period of improving soil-protective technology (1977-1986) increased 1.5 times in comparison with the previous period (1969-1976) and amounted to 3.3 t/ha (absolutely dry substance).

However, soil-protective technology did not guarantee an annual maturation to milky-wax ripeness. Only in this phase maize is the standard raw material (more than 25% of absolutely dry substance) for the production of high-quality silage [18].

There are not enough effective temperatures for late-maturing hybrids to reach the milky-wax ripeness until the autumn frosts in the climatic conditions of the Northern Kazakhstan.

The dynamics of the yield of maize in selected agricultural areas during this period and quality of the yield can be seen in a typical example of the North-Kazakhstan region. We analyzed the maize cultivation in the region by the equal squares method during the 17-year period.

As convincingly demonstrated by mathematical analysis (table 2), in all agricultural areas during the period of years 1969-1986 the transition to soil-protective technology has affected the trend of growth of the yields, expressed as a positive value of the parameter x (average annual factor).

Table 1 – The technological cycles in the different periods of cultivation of maize at the experimental field of the North-Kazakhstan Scientific and Research Institute of Agriculture

Technological operation	Technology	
	Tilled (1969-1976)	Soil-protective (1977-1986)
Ploughing	+	–
Treatment of the ground with a cultivator	–	+
Harrowing with spike-tooth harrows	+	–
Harrowing with needle harrows	–	+
Treatment of the ground by disk plough-harrow before sowing	+	–
Treatment of the ground with a cultivator before sowing	–	+
Packing soil before sowing	+	–
Checkrow planting	+	–
dotted sowing with fertilizers and herbicides	–	+
Packing soil after sowing	+	–
Double harrowing	++	++
Treatment of soil in row-spacing	+	+
Treatment of soil in row-spacing with introduction of fertilizers	+	–
Treatment of soil in row-spacing crosswise the direction of sowing, 2 passes	++	–
Treatment of plants with herbicides	+	–
Harvesting	+	+
The number of treatments	14	8
The average yield for the period, t/ha	2.2	3.3

Table 2 – The tendency in dynamics of the yield of maize in agricultural areas of the North-Kazakhstan region (during the period of years 1969-1986)

Agricultural zone	The tendency in dynamics of the yield	Yield		
		Natural humidity, t/ha	Absolutely dry substance	
			Concentration, %	Yield, t/ha
Moderately arid steppe	92.6 +4.4 x	14.9	20.4	3.03
Arid steppe	76.0 +5.5 x	13.2	23.0	3.03
Forest-steppe	112.0 +1.5 x	13.6	21.4	2.91
Hilly steppe	121.0 +2.7 x	15.6	19.5	3.04
Steppe on chernozem soils	105.0 +2.3 x	13.4	19.2	2.57
Steppe on chestnut soils	37.9 +2.2 x	7.09	20.0	1.41
Experimental field of The North-Kazakhstan Scientific and Research Institute of Agriculture (hilly steppe)	169.3 +5.7 x	22.6	23.1	5.22

Despite the fact that high technological discipline was maintained on the experimental field, as evidenced by the highest yield here, there was no guarantee for the milk-wax ripeness also on this field. The concentration of absolutely dry substance was lower than 25% that did not provide a producing of high-quality silage.

The concentration of absolutely dry substance in mid-season hybrids (FAO 200-250) which have been approved for use in the Northern Kazakhstan was reaching 20-23% before the first frost. The main cause of low-quality silage is the biomass with high humidity.

Since 1987, the team of scientists of the North-Kazakhstan Scientific and Research Institute of Agriculture started to develop an improved technology. This technology was based on the introduction of early-maturing hybrids (FAO 150-199). These hybrids had milky-wax ripeness every year in this region [19].

The yield of early-maturing hybrids was higher than that of the mid-season hybrids, as was proven by our research and confirmed by verification in production. It is well known that the maximum daily gain of maize falls on the phase of milky-wax ripeness and significantly exceeds the rates of all the previous phases [20-24].

Early maturing hybrids grow intensively 12-15 days until frost, while for the mid-season hybrids this period is 1-3 days and is interrupted by frost. Therefore, the mid-season hybrids which have a higher potential of productivity in comparison with the early-maturing hybrids, inferior to them in the yield of absolutely dry substance in the short summer.

The average yield for the 1987-1995 period was higher (advanced technology) than for the prior periods (figure 1).




Figure 1 – Dynamics of the yield of maize in the experimental field of the North-Kazakhstan Scientific and Research Institute of Agriculture as shown by the periods of cultivation

However, the improved technology does not guarantee the annual high yields in difficult climatic conditions of the Northern Kazakhstan. Maize was the main silo crop in the region, other silage crops occupied 6-17% of the total area.

Therefore, fluctuations in yield depended only on one culture. Maize harvest is increasing slowly in years with cool summers; the moisture content of the plants will be high. Maize also gives low yields in dry years (table 3).

To the main silo crop needs a set of other cultures for a particular agricultural zone. This set of other cultures will reduce the fluctuations of annual yield. Average yield of these crops should be higher than the yield of maize. In this case, we obtain the optimal set of crops [25].

The optimal period for harvesting of maize corresponds to the phase of milky-wax ripeness and lasts 8-10 days. In fact, this period lasts 18-20 days, if maize is a monoculture. Therefore, harvesting silage ends, usually at subzero temperatures, which affect the quality of biomass.

Table 3 – The amplitude of fluctuations of yield of maize in the experimental field of the North-Kazakhstan Scientific and Research Institute of Agriculture

Year	The sum of temperatures above +10°C, °C	The amount of precipitation (May-August), mm	The period without effective rainfall, days	Absolutely dry substance, t/ha	The decrease in yield, t/ha
1987	1775	132.2	30.VI – 7.VII (37)	3.0	1.8
1988	2120	129.3	27.VI – 10.VII (15)	5.7	–
1989	2178	133.1	27.VI – 21.VIII (51)	2.0	2.8
1990	2077	304.2	29.V – 15.VI (16)	7.3	–
1991	2124	132.2	31.V – 7.VII (37)	4.2	0.6
1992	1890	281.4	8.VI – 21.VI (12)	7.6	–
1993	1951	238.9	3.VI – 5.VII (32)	4.0	0.8
1994	1960	287.6	20.VI – 7.VII (16)	5.8	–
1995	1627	122.4	20.VI – 9.VII (19)	3.7	1.1
1996	1886	219.0	21.VI – 8.VII (16)	4.8	–
1997	2230	144.0	17.VI – 5.VII (18)	4.7	0.1
1987-1997	1983	194.5	(24)	4.8	

Note: Rainfall of more than 5 mm per day classified as effective.

Typical results were obtained by us in 1996, when the first frost was on August, 26 (table 4).

Table 4 – Chemical composition of maize in different periods of harvesting (1996)

Indicators of the quality of the yield	Unit of measurement	Period of harvesting		
		Milk ripeness	Milky-wax ripeness	
			Before the frost	After the frost
Date of sampling		20.VIII	25.VIII	30.VIII
Humidity	%	79.3	75.3	77.8
Absolute dry substance	%	20.7	24.7	22.2
Including				
protein	%	3.0	2.6	2.5
Fat	%	0.4	0.8	0.2
cellulose	%	7.4	6.9	11.5
nitrogen-free extractives	%	8.0	12.6	6.6
Ash	%	1.9	1.8	1.4
The amount of carotene	mg / kg	23.5	15.9	8.6

The researchers had to find the set of crops in the system of silage conveyor to decrease the amplitude of harvest variations over years with extreme conditions. At the same time maize should remain the main silo crop. The results of the research (1998-2006) showed that in years with cool summers, when the number of active temperatures is insufficient, sunflower gives the greatest yield. The yield of sugar sorghum is much greater than yield of maize and sunflower in dry land conditions when the period of vegetation is prolonged [25].

The yield of maize, sweet sorghum and sunflower differ from each other and depended on the period of harvesting (table 5).

Yield of sunflower, compared with yield maize and sorghum, had the maximum value on the 25 of August. Because the maize did not reach milk-wax ripeness yet, as well as sorghum did not reach the stage of flowering. Maize reached milky-wax ripeness on the 30 of August, and on this day, the yield of maize was higher than that of sunflower and sugar sorghum. In turn, sweet sorghum reached its maximum yield in the first five days of September.

Table 5 – The yield of silage crops by calendar dates (absolutely dry substance), t/ha

Year	Date of harvesting			
	August, 20	August, 25	August, 30	September, 5
Maize (control)				
2007	18.2	40.1	48.0	–
2008	30.2	62.3	68.4	–
2009	24.6	50.3	55.5	–
2007-2009	24.0	50.2	57.3	–
Sun flower				
2007	59.1	65.2	–	–
2008	38.3	45.0	–	–
2009	49.5	56.9	–	–
2007-2009	48.0	55.0	–	–
Sweet sorghum				
2007	19.3	24.5	26.3	38.5
2008	24.5	36.0	47.6	69.2
2009	22.4	35.2	44.7	62.4
2007-2009	21.0	31.2	39.6	56.7

The interpretation of these data allows to calculate the yield of silage crops in the system of silage conveyor, if the harvesting period fits: for sunflower - August, 20-25 (flowering stage); for maize - August, 26-30 (milky-wax ripeness), sweet sorghum - August, 31 – September, 5 (flowering of panicles).

We compared the average yield of crops in the system of silage conveyor (maize, sunflower and sugar sorghum) with a yield of monoculture of maize over 3 years (2007-2009). It was found (table 6) that the amplitude of fluctuation of yield for the monoculture of maize was higher from year to year. The average yield of monoculture (2007-2009) was lower, because the harvesting period was prolonged. Harvesting of monoculture of maize began before reaching the milk-wax ripeness.

Table 6 – Comparison of different systems for the production of silage

The system of the production	2007	2008	2009	2007-2009
Monoculture of maize, t/ha	3.54	5.36	4.34	4.41
The deviation from the mean value				
t/ha	- 0.87	+ 0.95	- 0.07	–
%	- 19.7	+ 21.5	- 1.5	
The system of silage conveyor, t/ha	4.61	5.51	5.32	5.14
The deviation from the mean value				
t/ha	- 5.53	0.37	0.18	
%	- 10.3	+ 7.1	+ 3.5	

The yield of crops in system of silage conveyor will be higher if corn, sunflower and sweet sorghum do (no “will” after “if”) not occupy equal areas. They should occupy the acreage according to the probability of occurrence of certain weather conditions.

It was established by us that the number of years with high solar activity and prolonged summer in the hilly-steppe zone was 8 of 30 (1961-1990). Therefore, the acreage of sweet sorghum should be no more than 25%, which corresponds to the theoretical frequency of coming of favorable years. The theoretical frequency of coming of favorable years for sunflower corresponds to 25-27%.

Conclusion. Scientific search has developed from tilled technology to soil-protective and further to advanced technology on the principles of the increasing yield, the quality of yield and minimizing of the technological cycle during the 5 decades of cultivation of maize in the Northern Kazakhstan.

The average yield was 4.8 t/ha (absolutely dry substance) over the years of the introduction of improved technology (1987-1997) at the experimental field of the North-Kazakhstan Scientific and Research Institute of Agriculture. Yields increased 2.2 times compared to the tilled technology (1969-1976). Labor costs decreased 4.9 times per 1 ton of product.

Silage conveyor was developed on the basis of the set of crops, which supplemented maize, for production of sustainable by years raw material silage. Yields of maize as a monoculture for production of silage amounted to 4.41 t/ha during 2007-2009, that is 0.73 t/ha less than the average yield of crops of the system of silage conveyor (sunflower - corn - sugar sorghum). The amplitude of fluctuation of yield decreased from 19.7% (monoculture) to 10.3% (silage conveyor).

REFERENCES

- [1] Goodman M.M. The history and origin of maize: current theories on the relationships between maize and some of its relatives. North Carolina Agricultural Experiment Station, 1965. 170 p.
- [2] Sazanova L. The history of distribution of maize in our country. Minsk: Urozhaj, 1964. 219 p.
- [3] Nalivaiko G.A. About the tilled farming system. The Publishing House of the Ministry of Agriculture of the RSFSR (Russian Soviet Federative Socialist Republic). Moscow, 1962. 112 p.
- [4] Berestovsky G.G. Wind erosion of soils and control measures against it in the north-east of the Kazakhstan: Abstract of dissertation of Ph.D. in agricultural sciences. Alma-Ata, 1969. 28 p.
- [5] Gossen E.F. Soil-protective farming and sustainability of grain production. Kokshetau: Poligrafija, 2014. 135 p.
- [6] Khlebov P.I. The characteristics of varietal farming of maize in the steppe zone of Northern Kazakhstan: dissertation of Ph.Ds. in agricultural sciences. Shortandy. 1966. 153 p.
- [7] Kozhevnikov A.R., Popova G.I. Maize in the Virgin Lands region and in the Western Siberia. M.: Kolos, 1965. 144 p.
- [8] Rabinovich N.M. Agro-technical and organizational-economic features of maize cultivation in Kazakhstan: Dissertation of Ph.Ds. in agricultural sciences. Alma-Ata, 1968. 103 p.
- [9] Erlepesov M.N. The cultivation of maize in Kazakhstan. The Publisher of agricultural literature, Alma-Ata, 1964. 228 p.
- [10] Krumin A.V. Maize in soil-protective system of farming // Maize. 1974. N 8. P. 14-15.
- [11] Zinchenko I.G. The influence of different systems of primary tillage on some elements of fertility and maize yield // Collection of scientific papers of The All-Union Scientific Research Institute of Grain Farming. 1970. P. 66-67.
- [12] Hall A.J., Lemcoff J.H., Trapani N. Water stress before and during flowering in maize and its effects on yield, its components, and their determinants. Maydica, 1981. P. 19-38.
- [13] Kudaybergenov G.K. The system of primary tillage in the grain crop rotation. Alma-Ata: Kainar, 1978. P. 14-16.
- [14] Pant M.M. Dependence of plant yield on density and planting pattern // Annals of Botany. 1979. N 44(4). P. 513-516.
- [15] Revut I.B. Theoretical foundations of elements of technological treatment of soil. Theoretical questions of treatment of soils. Leningrad, 1969. P. 47-49.
- [16] Bugutsky A.A. Use of the least squares method in studying the dynamics of the yield, cost value and productivity of labor // The Bulletin of Agricultural Science. 1963. N 6. P. 138-144.
- [17] Kostikov I.F. Combined furrow-opener: the patent of the Republic of Kazakhstan № 17020, was declared 08.09.2004.
- [18] GOST 23638-90 1991. Silage from the green plants. Technical requirements. Interstate standard, M. 7 p.
- [19] Murzhanov I.T, Kostikov I.F. Improvement of the technology of production of maize for silage. Proceedings of the Republican Scientific and Practical Conference. Kokshetau, 2000. P. 84-86.
- [20] Hrushka J.A. Monograph about the maize. M.: Kolos, 1965. 751 p.
- [21] Srinivasan G. International Maize and Wheat Improvement Center (CIMMYT) // Maize Production and Improvement in Central Asia and Caucasus. Mexico, 2000. 135 p.
- [22] Srinivasan G. Serving the needs of Resource – Poor Maize Farmers in the Developing World // Maize Production and Improvement in Central Asia and Caucasus. Almaty, 2000. P. 6-15.
- [23] Srinivasan, G. International Maize and Wheat Improvement Center (CIMMYT) // Maize Production and Improvement in Central Asia and Caucasus. Mexico, 2001. P. 6-10.
- [24] Yugenhaymer R.U. Maize: Improvement of varieties. Seed production. M., 1979. 518 p.
- [25] Tynykulov M.K. Improvement of methods of cultivation of sugar sorghum for silage in terms of hill-plain area of Northern Kazakhstan: Diss. kand. of agricultural Sciences. Almaty, 2010. 149 p.

Т. Қ. Салихов¹, М. Қ. Тынықұлов¹, И. Ф. Костиков²,
Ж. Ш. Ургалиев¹, Г. М. Салхожаева¹, Ш. Е. Арыстанова¹

¹Л. Н. Гумилев атындағы Еуразия ұлттық университеті, Астана, Қазақстан,

²Ш. Уәлиханов атындағы Көкшетау мемлекеттік университеті, Қазақстан

ЖҮГЕРІДЕН АСТЫҚ ПЕН СҮРЛЕМ ӨНДІРУ БИОЛОГИЯЛЫҚ ЕРЕКШЕЛІКТЕРІ

Аннотация. Солтүстік Қазақстанда жүгері өсірудің технологиясын жетілдіру топырақты қорғау және сүрлем дайындау принципте негізделеді. Ерте пісетін будандарды қолдана отырып (ФАО (Азық-түлік және Ауыл шаруашылығы Ұйымы) 150-199) технологиялық циклды азайту негізінде озық технология Солтүстік Қазақстанның ауыл шаруашылығы ғылыми-зерттеу институтында 1987-1997 жылдары әзірленді. Осы кезеңде абсолютті құрғақ заттардың орташа шығуы 4,8 т/га құрайды, 45,1%-ға алдындағы кезеңнен артық (1977–1986 жж.). Алайда, озық технологиялар сүрлемге өсіруге ғана (монодақыл) негізделген. Сондықтан, жазы қысқа, салқын және құрғақшылық жылдары өнімділіктің күрт төмендеуі де байқалуы мүмкін. Сүрлемдік конвейерде жүгері монодақылға альтернатива ретінде өсу жағдайларына сәйкес әртүрлі биологиялық талаптарымен дақылдардың таңдауы негізделген. Сүрлемдік конвейерде дақылдардың өнімділігі (күнбағыс-жүгері-шай жүгері) 5,14 т/га құрады (абсолютті құрғақ зат бойынша), ол 2007 жылдан 2009 жылға дейін кезеңіндегі құралған жүгері монодақыл өнімділігінен 0,73 т/га жоғары болды.

Түйін сөздер: жүгері, сүрлем, жемшөп, астық, ерте пісетін будан, құрғақ зат, өсіру технологиясы, өсіру әдістері, өнімділік, сүрлемдік конвейер, монодақыл, Солтүстік Қазақстан.

Т. Қ. Салихов¹, М. Қ. Тынықұлов¹, И. Ф. Костиков²,
Ж. Ш. Ургалиев¹, Г. М. Салхожаева¹, Ш. Е. Арыстанова¹

¹Евразийский национальный университет им. Л. Н. Гумилева, Астана, Казахстан,

²Кокшетауский государственный университет им. Ш. Уәлиханова, Казахстан

БИОЛОГИЧЕСКИЕ ОСОБЕННОСТИ ПРОИЗВОДСТВА ЗЕРНА И СИЛОСА ИЗ КУКУРУЗЫ

Аннотация. Совершенствование технологии выращивания кукурузы в Северном Казахстане основывается на принципах почвозащитного земледелия и заготовки силоса. Передовая технология с применением раннеспелых гибридов (ФАО (Продовольственная и Сельскохозяйственная Организация) 150-199) на основе минимизации технологического цикла была разработана в Северо-Казахстанской научно-исследовательского института сельского хозяйства в период с 1987 по 1997 год. Средний выход абсолютно сухого вещества за этот период составила 4,8 т/га, что составляет 45,1% больше, чем за предыдущий период (1977–1986). Однако передовые технологии основывались только на выращивании на силос (монокультура). Вследствие этого нельзя исключить резкое снижение урожайности в засушливые годы и в годы, когда лето короткое и прохладное. В силосном конвейере в качестве альтернативы в монокультуре кукурузы был обоснован выбор культур с различными биологическими требованиями к условиям роста. Урожайность культур в силосном конвейере (подсолнечник-кукуруза-сорго) была 5,14 т/га (абсолютно сухое вещество), что составляет 0,73 т/га выше урожайности монокультуры кукурузы в период с 2007 по 2009 год.

Ключевые слова: кукуруза, силос, фураж, зерно, раннеспелый гибрид, сухое вещество, технология выращивания, методы выращивания, урожайность, силосный конвейер, монокультура, Северный Казахстан.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://geolog-technical.kz/index.php/kz/>

Верстка *Д. Н. Калкабековой*

Подписано в печать 13.02.2018.
Формат 70x881/8. Бумага офсетная. Печать – ризограф.
12,0 п.л. Тираж 300. Заказ 1.