ҚАЗАҚСТАН РЕСПУБЛИКАСЫ ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

ХАБАРЛАРЫ

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК РЕСПУБЛИКИ КАЗАХСТАН Казахский национальный исследовательский технический университет им. К. И. Сатпаева

NEWS

OF THE ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN Kazakh national research technical university named after K. I. Satpayev

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

1 (433)

JANUARY – FEBRUARY 2019

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR



NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Етегдіпд Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Ехрапдед, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Webof Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Етегдіпд Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Бас редакторы

э. ғ. д., профессор, ҚР ҰҒА академигі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Редакция алкасы:

Абаканов Т.Д. проф. (Қазақстан)

Абишева З.С. проф., академик (Қазақстан)

Агабеков В.Е. академик (Беларусь)

Алиев Т. проф., академик (Әзірбайжан)

Бакиров А.Б. проф., (Кырғыстан)

Беспаев Х.А. проф. (Қазақстан)

Бишимбаев В.К. проф., академик (Қазақстан)

Буктуков Н.С. проф., академик (Қазақстан)

Булат А.Ф. проф., академик (Украина)

Ганиев И.Н. проф., академик (Тәжікстан)

Грэвис Р.М. проф. (АҚШ)

Ерғалиев Г.К. проф., академик (Қазақстан)

Жуков Н.М. проф. (Қазақстан)

Кожахметов С.М. проф., академик (Казахстан)

Конторович А.Э. проф., академик (Ресей)

Курскеев А.К. проф., академик (Қазақстан)

Курчавов А.М. проф., (Ресей)

Медеу А.Р. проф., академик (Қазақстан)

Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)

Нигматова С.А. проф. (Қазақстан)

Оздоев С.М. проф., академик (Қазақстан)

Постолатий В. проф., академик (Молдова)

Ракишев Б.Р. проф., академик (Қазақстан)

Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)

Сейтмуратова Э.Ю. проф., корр.-мушесі (Қазақстан)

Степанец В.Г. проф., (Германия)

Хамфери Дж.Д. проф. (АҚШ)

Штейнер М. проф. (Германия)

«КР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,

http://www.geolog-technical.kz/index.php/en/

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: К. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор

д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Редакционная коллегия:

Абаканов Т.Д. проф. (Казахстан)

Абишева З.С. проф., академик (Казахстан)

Агабеков В.Е. академик (Беларусь)

Алиев Т. проф., академик (Азербайджан)

Бакиров А.Б. проф., (Кыргызстан)

Беспаев Х.А. проф. (Казахстан)

Бишимбаев В.К. проф., академик (Казахстан)

Буктуков Н.С. проф., академик (Казахстан)

Булат А.Ф. проф., академик (Украина)

Ганиев И.Н. проф., академик (Таджикистан)

Грэвис Р.М. проф. (США)

Ергалиев Г.К. проф., академик (Казахстан)

Жуков Н.М. проф. (Казахстан)

Кожахметов С.М. проф., академик (Казахстан)

Конторович А.Э. проф., академик (Россия)

Курскеев А.К. проф., академик (Казахстан)

Курчавов А.М. проф., (Россия)

Медеу А.Р. проф., академик (Казахстан)

Мухамеджанов М.А. проф., чл.-корр. (Казахстан)

Нигматова С.А. проф. (Казахстан)

Оздоев С.М. проф., академик (Казахстан)

Постолатий В. проф., академик (Молдова)

Ракишев Б.Р. проф., академик (Казахстан)

Сеитов Н.С. проф., чл.-корр. (Казахстан)

Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)

Степанец В.Г. проф., (Германия)

Хамфери Дж.Д. проф. (США)

Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online).

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов

Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

http://nauka-nanrk.kz/geology-technical.kz

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

Editorial board:

Abakanov T.D. prof. (Kazakhstan)

Abisheva Z.S. prof., academician (Kazakhstan)

Agabekov V.Ye. academician (Belarus)

Aliyev T. prof., academician (Azerbaijan)

Bakirov A.B. prof., (Kyrgyzstan)

Bespayev Kh.A. prof. (Kazakhstan)

Bishimbayev V.K. prof., academician (Kazakhstan)

Buktukov N.S. prof., academician (Kazakhstan)

Bulat A.F. prof., academician (Ukraine)

Ganiyev I.N. prof., academician (Tadjikistan)

Gravis R.M. prof. (USA)

Yergaliev G.K. prof., academician (Kazakhstan)

Zhukov N.M. prof. (Kazakhstan)

Kozhakhmetov S.M. prof., academician (Kazakhstan)

Kontorovich A.Ye. prof., academician (Russia)

Kurskeyev A.K. prof., academician (Kazakhstan)

Kurchavov A.M. prof., (Russia)

Medeu A.R. prof., academician (Kazakhstan)

Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)

Nigmatova S.A. prof. (Kazakhstan)

Ozdoyev S.M. prof., academician (Kazakhstan)

Postolatii V. prof., academician (Moldova)

Rakishev B.R. prof., academician (Kazakhstan)

Seitov N.S. prof., corr. member. (Kazakhstan)

Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)

Stepanets V.G. prof., (Germany)

Humphery G.D. prof. (USA)

Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the

Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

http://nauka-nanrk.kz/geology-technical.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Geological Sciences named after K.I. Satpayev

69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 1, Number 433 (2019), 223 – 230

https://doi.org/10.32014/2019.2518-170X.27

UDC 663

M. V. Gernet¹, I. N. Gribkova¹, K.V. Kobelev¹, D.E. Nurmukhanbetova², E.K. Assembayeva²

¹All-Russian Research Institute of Brewing, Non-Alcoholic and Wine Industry - branch of the Gorbatov's Federal Research Center of Food Systems of RAS, Moscow, Russia

²Almaty Technological University, Almaty, Kazakhstan.

E-mail: institut-beer@mail.ru, k.kobelev55@mail.ru, dinar2080@mail.ru, elmiraasembaeva@mail.ru

BIOTECHNOLOGICAL ASPECTS OF FERMENTED DRINKS PRODUCTION ON VEGETABLE RAW MATERIALS

Abstract. Fermented drinks are considered as the optimal form of a food product, used to enrich the diet with biologically active substances that have a beneficial effect on the metabolism and immunoresistance of the human body, which are provided not only by extraction, but also are formed as a result of the vital activity of various types of microorganisms. The latter, in turn, produce vitamins, amino acids, organic acids, etc., which gives fermented drinks a higher biological value. Drinks technology with the use of tea fungus (*Medusomyces gisevii*) is oriented for home cooking and it is not possible to apply it in an industrial environment. The relevance of this research work is to develop the technology of the drink, enriched with biologically active substances in the most accessible form to expand the range of functional beverages aimed at maintaining the health of various population groups. In order to solve this problem a combined yeast and lactic acid bacteria ferment was chosen, the optimal composition of the nutrient medium based on green tea and barley malt wort was determined, and the amino acid and vitamin activator was selected to reduce the fermentation process. The developed technology made it possible to obtain a drink with mixed leaven, harmonious in organoleptic characteristics, with a high content of amino acids, especially essential ones, which made it possible to recommend this functional drink for various groups of the population.

Keywords: tea, wort, microorganism cultures, yeast food, secondary fermentation products.

Introduction. Drinks are rightfully considered to be the most rational form of a food product for enriching the diet with biologically active substances and creating special-purpose products [1, 2]. The high popularity of beverages among the population suggests the constant development of technical, methodological, technological bases, including in the framework of the fight against fraud [3-7]. In recent years, the problem of their excessive consumption is actualized [8].

One of the priorities of the state policy of the Russian Federation in the field of healthy nutrition is the development of fermentation drink technologies, based on the use of plant raw materials and microorganism cultures, that produce a complex of substances with significant potential of useful properties [1]. According to many scientists, the use of fermentation drinks contributes to the correct metabolism and strengthens the immunoresistance of the human body, which are formed as a result of microorganism vital activity [1, 2, 9-11, 26].

One of the urgent tasks in the production of fermented drinks is the search for fermentation microorganisms, producing biologically active substances, which are necessary for the normal functioning of mankind during their life. The latter are most fully accumulated in the case of the symbiotic cultures use, which can be a single organism (for example, *Medusomyces gisevii*) or a cultures combination (for example, yeast and lactic acid bacteria) [1].

Literature review. The drink, obtained with use of *Medusomyces gisevii*, which consists of yeast cells and acetic acid bacteria, contains glucuronic, folic and citric acids, vitamins B_1 , B_2 , B_3 , B_6 , B_{12} , as well as difficult to determine substances with antibiotic action, including medusin. The viability of using

bacterial cellulose, obtained from *Medusomyces gisevii* as a hemostatic agent in veterinary medicine and humane medicine was confirmed [12]. A necessary condition for the life of this microorganism is the presence of tea in the raw materials composition.

Analysis of scientific and technical information suggests, that the creation of industrial technology, based on the use of cultures of microorganisms with known properties, that allow to control the process of drink production, is promising and relevant. At the same time, the conditions for obtaining a homemade drink described in the literature and patents for a method for producing a drink, based on tea fungus and attempts to create analogues with *Medusomyces gisevii* have not been continued [13-16].

Methods. To assess the effectiveness of technological methods, research methods used in the industry were used: determination of the volume fraction of alcohol - GOST 12787-81, the arbitration method, which is the separation of the alcohol fraction of a drink from extractive substances by distillation [17]; determination of titratable acidity - GOST 12788-87, which is a method for the titration of the acid content by neutralizing them with a 0.1 sodium hydroxide solution [18]; determination of the mass concentration of volatile components - a gas chromatography method with the Kristall 5000.1 flame ionization detector ("Khromatek", Russia) equipped with an automatic information collection and processing system [19]; determination of the mass concentration of organic acids – high-performance liquid chromatography with an "Agilent Technologies 1200" ("Agilent", USA) diode array detector [20]; determination of the mass concentration of sugars and glycerol - a method of high-performance liquid chromatography with a refractometric detector "Agilent Technologies 1200" ("Agilent", USA), equipped with an automatic system for collecting and processing information [21]; determination of the mass concentration of amino acids is a high performance liquid chromatography method with an "Agilent Technologies 1200" ("Agilent", USA) diode array detector, equipped with an automatic information collection and processing system [22].

The replication of experiments at all stages of the experiment - not less than 3. The results of experimental studies were processed by methods of mathematical statistics using Student's criterion.

In order to conduct a study on the production of fermented drinks, the following materials were used: barley malting brewing malt; brewing tricyclic malt light; black tea and green leaf; mixed ferment on the basis of dry brewing yeast S. cerevisiae and lactic acid bacteria *Betabacterium breve*.

Results. As part of the study, the following tasks were set: to investigate the possibility of using various plant raw materials and mixed cultures of microorganisms traditional for the brewing and non-alcoholic industries in the technology of fermented drink; select the optimal fermentation conditions; get a drink with stable performance over a long shelf life without loss of compounds, responsible for the functional orientation of the finished drink.

To identify the optimal composition of the nutrient medium, allowing to obtain a harmonious drink from the point of organoleptic and physicochemical parameters view, provided that yeast and lactobacilli are used, the following plant materials were used in the study: leaf tea (black and green), barley based wort and tritium-potassium malt brewing, as well as granulated sugar, in various ratios, and the fermentation was carried out by making mixed yeast brewer's yeast and lactic acid bacteria d at the rate of 4% by volume of the nutrient medium.

During the experiments, the duration of fermentation, controlled by loss of dry substances, using black and green tea coincided and amounted to 141 hours with barley malt wort and 114 hours with triticali wort, which is explained by the rich amino-acid composition of triticali malt wort, which accelerates enzymatic processes associated with the activity of yeast. However, the organoleptic evaluation of all analyzed samples using a 25-point system showed, that the most balanced indicators were fermented drink using a carbohydrate-tea base (based on green tea) and barley malt wort, in particular, a variant of the nutrient medium, based on green tea and wort from barley malt in a ratio of 75% and 25%, the data are presented in table 1.

In all other samples, the alcohol content and acidity were at 5–10% higher than the optimal sample, but an autolysis tone was present, which is unacceptable.

However, the duration of cultivation was quite long, so the next stage of our work was the use of various activators to shorten the process. For this purpose, the most productive amino acid-vitamin activator (AVA), obtained according to the technology [23] and containing 40–42% of soluble protein, including 13–15% of amine nitrogen, and vitamins (thiamine (B_1) , pantothenic acid (B_3) , pyridoxine (B_6) , biotin (B_7) , inoside (B_8) - 17 mg/kg. The share of AVA was 2.5–10 ml/hl or 0.5–2% by volume of the fermented wort (table 2).

Indicators

Characteristics of Fermented Green Tea and Barley Malt Wort

Organoleptic Indicators

Taste

fermented drink with slightly sour

Colour

straw

Aroma

pleasant aroma of a well fermented fruit drink

Physicochemical Indicators

Alcohol content, vol.%

3,30±0,2

Acidity, a.u.

11,00±0,3

Table 1 – Organoleptic and physicochemical indicators of the fermented wort optimal variant

Table 2 – Applied feedings doses for fermentation

	Nutrient medium variant						
Indicator name	Control	with Feedings of AVA, ±0,05					
	I	II	III	IV	V		
Applied feedings doses, %	_	0,5	1	1,5	2,0		

The dose of dressing with the use of selected nutrient medium (75% of tea and 25% of barley malt wort) is presented in table 2, the dynamics of acidity accumulation during fermentation - in figure 1, the content of secondary fermentation products – in the table 3. It should be noted that we previously studied other supplements, but they did not show any tangible effect [24].

Table 3 – The Content of Organic Acids (OA), Sugars and Glycerin in the Drink

Component	Content, mg/dm ³	Component	Content, mg/dm ³	Component	Content, mg/dm ³	
OA±5,0%	**	Amino Acids ± 5,0%**				
Oxalic acid	0,070	Aspartic	27,0	Tryptophan*	25,5	
Wine acid	0,104	Glutamic	29,2	Isoleucine*	36,0	
Malic acid	0,303	Asparagine	36,8	Phenylalanine*	17,5	
Lactic acid	4,614	histidine*	20,0	Leucine*	65,0	
Citric acid	0,147	Serine	48,0	Lysine*	5,0	
Succinic acid	0,257	Glutamine	22,0	Alanine	22,0	
The amount of volatile acids, in terms of acetic	0,18±2%	Arginine*	17,5	Tyrosine	16,8	
		Threonine*	126,5	Valine*	41,0	
		Glycine	40,0	Methionine*	49,0	
Sugars and Glycerin ± 4,0%**						
Fructose	10,480	Glucose	5,310	Sucrose	2,7000	
* - irreplaceable amino acids; ** - method reproducibility.						

The data in table 3 show, that the resulting drink had a wide range of functional compounds, responsible for functional orientation, and more significant accumulation of lactic acid occurs, as compared with the control, since the lactic acid bacteria Betabacter breve was used during fermentation. Sugar and glycerin in the finished drink is formed less than in the control, which is explained by a more intensive activity of microorganisms that consume sugar for reproduction. The composition of amino acids are also essential in 2 times more, than the control, which is explained by the most active metabolism of microorganisms.




Figure 1 – Dynamics of acidity accumulation in the samples when adding feedings to the nutrient medium during fermentation

Comparative studies have been carried out on fermented drink with use of tea fungus and mixed teabased leaven.

The composition of volatile compounds and sugars are presented in tables 4, 5. It should be noted that the higher alcohols in the resulting drink, compared with the drink on the tea fungus, accumulate much more (except for l-propanol), and the content of esters is not so different.

Table 4 – Content of volatile compounds, organic acids and sugars in drink with use of the microorganism medusomyces gisevii

Component	Content mg/dm ³	Component	Content mg/dm ³	Component	Content mg/dm ³	
Ethers, $mg/dm^3 \pm 10\%$ **	Amino Acids ± 5,0%**					
Ethyl acetate	2,89	Aspartic	10,9	Valine*	1,44	
Ethyl lactate	5,79	Glutamic	1,8	Methionine*	_	
Organic Acids, $g/dm^3 \pm 5.0\%$ **		Аспарагин	Asparagine	Трипто-фан*	Tryptophan*	
Oxalic acid	_	histidine*	0,8	Isoleucine*	_	
Wine acid	_	Serine	1,6	Phenylalanine*	2,0	
Malic acid	0,021	Glutamine	-	Leucine*	0,9	
Lactic acid	0,059	Arginine*	1,0	Lysine*	0,8	
Citric acid	0,0083	Glycine	1,2	Tyrosine	_	
Succinic acid	0,061	Threonine*	4,5	Totally Irreplaceable	11,4	
		Alanine	1,3			
Sugars, $mg/cm^3 \pm 4.0\%**$		•				
Fructose	3,39	Glucose	2,74	Sucrose	17,33	

According to the content of amino acids, more intensive accumulation occurs in the drink with use of mixed yeast leaven and lactic acid bacteria, including irreplaceable bacteria, accumulates 36 times more compared to the drink on the tea fungus.

Component	Content, mg/dm ³	Component	Content, mg/dm ³	Component	Content, mg/dm ³			
Ethers, $mg/dm^3 \pm 10\%$ **			Amino Acids ± 5,0%**					
Ethyl acetate	1,3	Aspartic	27,0	Valine*	41,0			
Ethyl lactate	3,3	Glutamic	29,2	Methionine*	49,0			
Organic Acids, $g/dm^3 \pm 5.0\%$	Organic Acids, g/dm ³ ± 5,0%**		36,8	Tryptophan*	25,5			
Oxalic acid	histidine*	histidine*	20,0	Isoleucine*	36,0			
Wine acid	0,373	Serine	48,0	Phenylalanine*	17,5			
Malic acid	4,83	Glutamine	22,0	Leucine*	65,0			
Lactic acid	0,059	Arginine*	17,5	Lysine*	5,0			
Citric acid	0,12	Glycine	40,0	Tyrosine	16,8			
Succinic acid	0,331	Threonine*	126,5	Totally Irreplaceable	403			
		Alanine	1,3					
Sugars, $mg/cm^3 \pm 4.0\%$ **	Sugars, mg/cm ³ $\pm 4.0\%$ **							
Fructose	19,9	Glucose	11,97	Sucrose	2,36			
* - irreplaceable amino acids; ** - method reproducibility.								

Table 5 – Content of volatile compounds, organic acids and sugars in the drink with use of mixed yeast leaven and lactic acid bacteria

Particular attention should be paid to the threonine amino acid, which accumulates 3 times more than in the drink with tea fungus (table 4, 5). From the point of view of the functional load, threonine is very useful for the body, since it is responsible for the protein metabolism in the body, is part of collagen, elastin and tooth enamel protein, positively affects the person's psycho-emotional state [25]. We assume that this drink, obtained by us with the help of a mixed leaven, can be recommended for a wide range of the population of Russia.

The third task to get a drink with stable performance over a long shelf life without losing the compounds, responsible for the functional orientation of the finished drink was solved using pasteurization techniques at a maximum processing temperature of 72 °C, due to which it was possible to extend the shelf life of the drink for 6 months.

Analysis of the secondary fermentation products in mixed leaven in a drink showed, that their concentration differs from that in a drink made on the basis of a tea fungus, which is explained by a different metabolic set of enzymes in two cases.

On the basis of the data, obtained in the study, the optimal organoleptic and physicochemical indicators of fermented drinks, prepared from pasteurized fermented bases were determined, which made it possible to develop a draft regulatory documentation for fermented drinks on plant raw materials.

Conclusions. Studies have allowed picking up planting materials and microbiological cultures, working out the technological parameters of the process, including fermentation modes - the optimal composition for the drink technology was the ratio of green tea extract and barley malt wort 3:1, as well as the use of an additive ava in the amount of 1%, which made it possible to reduce the fermentation time to 93 hours and to obtain a drink with high organoleptic and physicochemical parameters.

On the basis of a comparison of the drink we developed, with based on tea fungus drink, at the main functional components, it was concluded that the content of amino acids was 4 times higher than the total amount and 3 times threonine, which is probably the basis recommend a drink for food of various groups of the population of the Russian Federation.

Discussion. The conducted literary search on the studied problem revealed a diverse composition of vegetable raw materials in order to apply it in the drink technology. This type of raw material is traditionally used in fermented drinks, based on the microorganism *Medusomyces gisevii* [1-3, 18]. However, this microorganism did not find wide application in industry due to insufficient knowledge of catabolic reactions.

Therefore, we were faced with the task of developing a technology for a drink, that would be an analogue of the "Kombucha" drink. For this purpose, were used yeast and lactic acid bacteria, traditionally used in the production of kvass.

The use of plant materials, in particular tea, and barley malt wort and mixed microbiological cultures, made it possible to obtain a fermented drink enriched with functional compounds such as organic acids, amino acids, ethers and other volatile compounds in quantities exceeding drinks with use of tea fungus culture. This fact is explained by the difference in the metabolism and the set of enzymes of the two microorganism cultures, since the in technology were used lactic acid bacteria together with yeast, which is characterized by its own set of secondary products of metabolism compared to the culture of *Medusomyces gisevii* [19].

The obtained data on the composition and content of alcohols in the drink confirm the correctness of the choice of the malting wort as a source of nitrogenous nutrition for yeast, since it is known that it is the amino acids, that participate in the formation of higher alcohols as a result of catabolic reactions. The use of yeast and lactic acid bacteria, as well as a balanced plant substrate (tea and malt wort) made it possible to obtain a harmonious drink, rich in functional compounds. Accelerated fermentation metabolism was promoted by the use of an amino acid activator in the above-mentioned concentration, since the activator influenced the growth and reproduction of yeast. This is of great importance, since the duration of fermentation has decreased to 93 hours. The use of pasteurization techniques has made it possible to increase the shelf life without loss of useful compounds.

This fact allows to introduce this technology in the enterprises of the beer and non-alcoholic industry, since the microorganisms used are well studied and their livelihoods can be regulated, that is, they are technological.

М. В. Гернет¹, И. Н. Грибкова¹, К. В. Кобелев¹, Д. Е. Нурмуханбетова², Э. К. Асембаева²

¹Бүкілресейлік сыра қайнату, алкогольсіз және шарап өнеркәсібі ғылыми-зерттеу институты — В. М. Горбатов атындағы «Азық-түлік өнімдерінің федералдық ғылыми орталығы» Федералдық мемлекеттік бюджеттің ғылыми мекемесінің филиалы РҒА, Мәскеу, Ресей,

²Алматы технологиялық университеті, Алматы, Қазақстан

ӨСІМДІК ШИКІЗАТЫН ҚОЛДАНЫП АШЫТЫЛАТЫН СУСЫНДАР ӨНДІРІСІНІҢ БИОТЕХНОЛОГИЯЛЫҚ АСПЕКТІЛЕРІ

Аннотация. Ферменттелген сусындар биологиялық белсенді заттармен тағам рационын байыту үшін пайдаланылатын тамақ өнімдерінің оңтайлы түрі болып табылады, олар адам ағзасының зат алмасуына және иммундык тұрақтылығына оң әсерін тигізеді, олар экстракция арқылы ғана емес, сондай-ақ түрлі микроорганизмдердің өмірлік белсенділігі нәтижесінде пайда болады. Олар өз кезегінде витаминдерді, аминқышқылдарын, органикалық қышқылдар және тағы басқа заттарды түзеді, бұл ферменттелген сусындардың биологиялық құндылығын арттырады. Шай саңырауқұлағы (Medusomyces gisevii) қолданылған сусындар технологиясы үйде алуға арналған және оны өнеркәсіптік жағдайда қолдану мүмкін емес. Осы ғылымизерттеу жұмысының өзектілігі әртүрлі халық топтарының денсаулығын сақтауға бағытталған функционалдық сусындардың ауқымын кеңейту үшін барынша қолжетімді түрде биологиялық белсенді заттармен байытылған сусынның технологиясын жасау болып табылады. Бұл мәселені шешу үшін аралас ашытқылар мен сүт қышқылдары бактерияларының ұйытқысы таңдап алынды, көк шай және арпа уыты негізіндегі қоректік ортаның оңтайлы құрамы анықталды, аминқышқылы мен витаминді активатор ашыту үрдісін азайту үшін таңдалды. Алынған технология құрамында ауыстырылмайтын амин қышқылдарының көп мөлшері бар, органолептикалық көрсеткіштері үйлесімді аралас ашытқы негізіндегі сусын алуға мүмкіндік берді, бұл халықтың түрлі топтары үшін осы функционалдық сусындарды ұсынуға мүмкіндік берді.

Түйін сөздер: шай саңырауқұлағы, арпа уыты, ашытқылар мен сүтқышқылды бактериялардың біріктірілген ұйытқысы.

М. В. Гернет¹, И. Н. Грибкова¹, К. В. Кобелев¹, Д. Е. Нурмуханбетова², Э. К. Асембаева²

¹Всероссийский научно-исследовательский институт пивоваренной, безалкогольной и винодельческой промышленности – филиал Федерального государственного бюджетного научного учреждения «Федеральный научный центр пищевых систем им. В. М. Горбатова» РАН, Москва, Россия,

²Алматинский технологический университет, Алматы, Казахстан

БИОТЕХНОЛОГИЧЕСКИЕ АСПЕКТЫ ПРОИЗВОДСТВА НАПИТКОВ БРОЖЕНИЯ С ПРИМЕНЕНИЕМ РАСТИТЕЛЬНОГО СЫРЬЯ

Аннотация. Напитки брожения рассматриваются как оптимальная форма пищевого продукта, используемого для обогащения рациона питания биологически-активными веществами, благотворно влияющими на обмен веществ и иммунорезистентность организма человека, которые обеспечиваются не только за счет экстракции, но и формируются в результате жизнедеятельности различных видов микроорганизмов. Последние, в свою очередь, продуцируют витамины, аминокислоты, органические кислоты и др., что и придает ферментированным напиткам повышенную биологическую ценность. Технология напитков с применением чайного гриба (Medusomyces gisevii) предназначена для домашнего приготовления и не представляется возможным применить ее в промышленных условиях. Актуальность данной исследовательской работы заключается в разработке технологии напитка, обогащенного биологически активными веществами в наиболее доступной форме для расширения ассортимента функциональных напитков, направленных на поддержание здоровья различных групп населения. С целью решения данной задачи была выбрана комбинированная закваска дрожжей и молочнокислых бактерий, определен оптимальный состав питательной среды на основе зеленого чая и ячменного солодового сусла, для сокращения процесса брожения был подобран аминокислотно-витаминный активатор. Разработанная технология позволила получить напиток на смешанной закваске, гармоничный по органолептическим показателям, с повышенным содержанием аминокислот, особенно незаменимых, что позволило рекомендовать этот функциональный напиток для различных групп населения.

Ключевые слова: чайный гриб, сусло, комбинированная закваска дрожжей и молочнокислых бактерий.

Information about authors:

Gernet M. V., Head of Brewring department, Professor, Doctor of technical Science, All-Russian Research Institute of Brewing Non-Alcoholic and Wine Industry - branch of the Gorbatov's Federal Research Center of Food Systems of RAS, Moscow, Russia; institut-beer@mail.ru; http://orcid.org/0000-0003-0277-7318.

Gribkova I. N., Researcher of Brewring department, Candidate of technical Sciences, All-Russian Research Institute of Brewing, Non-Alcoholic and Wine Industry - branch of the Gorbatov's Federal Research Center of Food Systems of RAS, Moscow, Russia; institut-beer@mail.ru; http://orcid.org/0000-0002-4373-5387.

Kobelev K. V., Deputy directore of Scintific research institut of Brewring, Candidate of technical Sciences, All-Russian Research Institute of Brewing, Non-Alcoholic and Wine Industry - branch of the Gorbatov's Federal Research Center of Food Systems of RAS, Moscow, Russia; k.kobelev55@mail.ru; http://orcid.org/0000-0002-3619-6282.

Nurmukhanbetova Dinara Erikovna, candidate of engineering sciences, acting associate professor, Almaty Technological University, Department of Food safety and quality; dinar2080@mail.ru; http://orcid.org/0000-0002-8939-6325.

Assembayeva Elmira Kuandykovna, Almaty Technological University, Department of Food Biotechnology, Master of Technical Sciences, Senior Lecturer; elmiraasembaeva@mail.ru; http://orcid.org/0000-0001-7964-7736.

REFEREVCES

- [1] Shenderov B.A., Doronin A.F. (2001). Perspectives of Functional Drinks for Various Population Groups on the Tea and Coffee Basis. Materials of the 1st International Specialized Exhibition "Tea and Coffee Magic Aroma". Moscow. Russian Exhibition Center (VVC). P. 58-60 (in Rus.).
- [2] Pilipenko T.V., Korotysheva L.B. (2016). Study of Tea-Based Drinks Quality and Functional Properties, South Ural State University Bulletin. Series "Food and Biotechnology", 4: 1. P. 87- 94 (in Rus.).
- [3] Khanferyan R.A., Vybornaya K.V., Rajabkadiev R.M. (2017). Frequency of consumption of sugary carbonated drinks by the population of different age groups of the Russian Federation. Nutrition Issues, 3. P. 55-58 (in Rus.).
- [4] Bessonov V.V., Khanferyan R.A., Galstyan A.G. (2017). Potential Side Effects of Caffeine Consumption in Healthy Adults, Pregnant Women, Adolescents and Children (A Review of Foreign Information). Nutrition Issues, 6. P. 21-28 (in Rus.).

- [5] Khanferyan R.A., Rajabkadiev R.M., Evstratova V.S. et al. (2018). Consumption of Carbohydrate-Containing Drinks and their Contribution to the Total Caloric Intake. Nutrition Issues, 87: 2. P. 39-43 (in Rus.).
- [6] Oganesyants L.A., Panasyuk A.L., Kuzmina E.I. (2016). Determination of the Carbon Isotope 13C/12C in Ethanol of Fruit Wines in Order to Identification Characteristics, Foods and Raw Materials, 4:1. P. 141-147. DOI: 10.21179/2308-4057-2016-1-141-147 (in Eng.).
- [7] Zyakun A.M. et al. (2012). Mass Spectrometry Analysis of Prevalence of 13C/12C Isotopes Ratios in Grape Plants and Wine in Dependence on Climatic Factors (Krasnodar Territory and Rostov Region, Russia), Mass Spectrometry, 9:2. P. 16-22. DOI: 10.1134/S106193481313011X (in Eng.).
- [8] Petrov A.N., Khanferyan R.A., Galstyan A.G. (2016). Current Aspects of Counteraction of Foodstuff's Falsification, Nutrition Issues, 85:5. P. 86-92 (in Rus.).
- [9] Sergeeva I.Yu., Unschikova T.A., Rysina V.Yu. (2014). Improvement Directions of Kvass Fermentation Technology Based on the Analysis of Modern Scientific and Technical Developments, Equipment and Technology of Food Production, 3. P. 69-78 (in Rus.).
- [10] Budaeva V.V., Gladysheva E.K., Skiba E.A., Sokovich E.V. (2016). The Method of Obtaining Bacterial Cellulose [Metod poluchenija bakterial'noj celulozi]. Patent of Russian Federation [Patent Rosijskoj Federacii] (in Rus.).
 - [11] Agbo F., Spradlin J.E. (1995). Enzymatic Clarification of Tea Extracts. Patent of USA (in Eng.).
 - [12] Khachatryan V. (2012). Tea Fungus: Sober Way Out. ISBN 978-5-88503-985-7. P. 99-101 (in Rus.).
 - [13] Shkitina E.N. (2012). Healing Fungi from All Diseases. ISBN 978-5-386-03870-0. P. 32-38 (in Rus.).
- [14] Khachatryan V.Kh., Ivanova T.V. (2000). Drink, Method of Tea Fungus Liquid Culture Production and Method of Drink Production [Napitki. Metod proizvodstva I poluchenia kulturi chajnogo griba]. Patent of Russion Federation [Patent Rosujskoj Federacii] (in Rus.).
- [15] GOST 12787-81 Beer. Methods for Alcohol Determination, Actual Extract and Calculation of Dry Substances in the Initial Wort (in Rus.).
 - [16] GOST 12788-87 Beer. Methods for Acidity Determining [GOST. Pivo. Obschie tehnicheskie uslovija] (in Rus.).
- [17] Methods for Measuring the Mass Concentration of Volatile Components in Fermented Products by Gas Chromatography (Certificate of Measurement Method Attestation no. 01.00225/205-45-11) [Hromatograficheskie metodi izmerenija masovoj koncentracii letuchih komponentov] (in Rus.).
- [18] Methods of Measuring the Mass Concentration of Organic Acids in Fermented Products by High Performance Liquid Chromatography (Certificate of Measurement Method Attestation no. 01.00225/205-49-12) [Hromatograficheskie metodi izmerenija masovoj koncentracii organiceskih kislot metodom visokoafektivnoj zhidkostnoj hromatografii] (in Rus.).
- [19] Methods for Measuring the Mass Concentration of Sugars and Glycerin in Alcoholic and Non-Alcoholic Drinks by High Performance Liquid Chromatography (Certificate of Measurement Method Attestation no. 01.00225/205-54-12) [Hromatograficheskie metodi izmerenija masovoj koncentracii saharov I glicerina v alkogolnih I bezalkogolnih napitkah metodom visokoafektivnoj zhidkostnoj hromatografii] (in Rus.).
- [20] Methods for Measuring the Mass Concentration of Amino Acids in Alcoholic and Non-Alcoholic Drinks by High Performance Liquid Chromatography (Certificate of Measurement Method Attestation no. 01.00225/205-48-12) [Hromatograficheskie metodi izmerenija masovoj koncentracii aminokislot v alkogolnih I bezalkogolnih napitkah metodom visokoafektivnoj zhidkostnoj hromatografii] (in Rus.).
- [21] Shaburova L.N., Ilyashenko N.G., Sadova A.I. [et al.] (2000). Yeast Activation Method [Metod aktivacii drozhej]. Patent of Russion Federation [Patent Rosijskoj Federacii] (in Rus.).
- [22] Gernet M.V., Kobelev K.V., Gribkova I.N., B.R. (2016). Development of Functional Fermented DrinksTechnology with Use of Tea. Part II. Formation of Main and By-Products of Fermentation during the Cultivation of Microorganisms [Issledovanie tehnologii funkcional'nih napitkov brozhenija na osnove chaja. Chast II. Obrazovanie osnovnih pobochnih produktov brozhenija v hode kul'tivirovanija mikroorganizmov]. Beer and Bevereges [Pivo I napitki], 2. P. 12-16.
 - [23] Shterman S.V. (2017). Sports Nutrition Products Moscow, Moscow. ISBN 978-5-906955-07-4(in Rus.).
- [24] Fu C., Yan F., Cao Z. et al. (2014). Antioxidant Activities of Kombucha, Prepared from Three Different Substrates and Changes in Content of Probiotics during Storage, 34:1. P. 123-126. DOI.org/10.1590/S0101-20612014005000012 (in Eng.).
- [25] Jayabalan R., Malini K., Sathishkumar M. (2010). Biochemical Characteristics of Tea Fungus, Produced during Kombucha Fermentation, Food Sci. Biotechnology, 19:3. P. 843-847. DOI/10.1007/s10068-010 (in Eng.).
- [26] Oganesyants L.A., Khurshudyan S.A., Galstyan A.G., Semipyatny V.K., Ryabova A.E., Vafin R.R., Nurmukhanbetova D.E., Assembayeva E.K. (2018). Base matrices invariant digital identifiers of food products News «Series of Geology and Technical Sciences». N 6. P. 6-15. ISSN 2224-5278 https://doi.org/10.32014/2018.2518-170X.30.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see http://www.elsevier.com/publishingethics and http://www.elsevier.com/journal-authors/ethics.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis as electronic preprint, or an see http://www.elsevier.com/postingpolicy), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyrightholder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service http://www.elsevier.com/editors/plagdetect.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

http://www.geolog-technical.kz/index.php/en/

Верстка Д. Н. Калкабековой

Подписано в печать 06.02.2019. Формат 70х881/8. Бумага офсетная. Печать – ризограф. 16,7 п.л. Тираж 300. Заказ 1.