

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының Ғылым Академиясының
Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Kazakh national research technical university
named after K. I. Satpayev

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

6 (438)

NOVEMBER – DECEMBER 2019

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Б а с р е д а к т о р ы
э. ғ. д., профессор, ҚР ҰҒА академигі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Р е д а к ц и я а л қ а с ы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., академик (Қазақстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қырғыстан)
Беспәев Х.А. проф. (Қазақстан)
Бишимбаев В.К. проф., академик (Қазақстан)
Буктуков Н.С. проф., академик (Қазақстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Ерғалиев Г.К. проф., академик (Қазақстан)
Жуков Н.М. проф. (Қазақстан)
Қожахметов С.М. проф., академик (Қазақстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., академик (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Нигматова С.А. проф. (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Қазақстан)
Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)
Сейтмуратова Э.Ю. проф., корр.-мүшесі (Қазақстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«ҚР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н н а я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., академик (Казахстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Беспаяев Х.А. проф. (Казахстан)
Бишимбаев В.К. проф., академик (Казахстан)
Буктуков Н.С. проф., академик (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Ергалиев Г.К. проф., академик (Казахстан)
Жуков Н.М. проф. (Казахстан)
Кожаметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., академик (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Нигматова С.А. проф. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Казахстан)
Сейтов Н.С. проф., чл.-корр. (Казахстан)
Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

E d i t o r i a l b o a r d:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., academician (Kazakhstan)
Agabekov V.Ye. academician (Belarus)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Bespayev Kh.A. prof. (Kazakhstan)
Bishimbayev V.K. prof., academician (Kazakhstan)
Buktukov N.S. prof., academician (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadjikistan)
Gravis R.M. prof. (USA)
Yergaliev G.K. prof., academician (Kazakhstan)
Zhukov N.M. prof. (Kazakhstan)
Kozhakhmetov S.M. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., academician (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Nigmatova S.A. prof. (Kazakhstan)
Ozdoev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Rakishev B.R. prof., academician (Kazakhstan)
Seitov N.S. prof., corr. member. (Kazakhstan)
Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 6, Number 438 (2019), 6 – 11

<https://doi.org/10.32014/2019.2518-170X.150>

Aang Panji Permana^{1,2}, Subagyo Pramumijoyo², Akmaluddin²

¹Gorontalo State University, Gorontalo, Indonesia,

²Universitas Gadjah Mada, Yogyakarta, Indonesia.

E-mail: aang@ung.ac.id, bagyo@ugm.ac.id, akmaluddin@gmail.com

UPLIFT RATE OF GORONTALO LIMESTONE (INDONESIA) BASED ON BIOSTRATIGRAPHY ANALYSIS

Abstract. The uplift rate of Gorontalo limestone is highly dependent on the completeness of age and depth data. All data needed can be obtained based on complete biostratigraphy analysis. The research material used was a 24 meter thick limestone out crop. The aim of the research was to determine the absolute age of limestone, paleobathimetry and uplift rate of limestone in the research area. The three methods used consisting of the measured section (MS), biostratigraphy analysis and tectonic analysis. Bio datum in the limestone is only one, namely LO *Globoquadrina dehiscens*. Determination of paleobathimetry using two methods. The uplift rate of limestone is 0.0699-0.0724 mm/year.

Keywords: uplift rate, limestone, Gorontalo, biostratigraphy.

Introduction. The Indonesian archipelago is geologically the center of the meeting of three of world's main active plates, the Indian-Australian Plate which moves north-northeast, the Pacific Ocean Plate moves west-northwest and the Eurasian Continent Plate is almost static [1-4].

Sulawesi Island, which is located in the central part of the Indonesian Archipelago resembles the letter K. This form is influenced by the movement of the three main plates of the world. Based on regional physiography, Sulawesi Island is divided into South Arm, Middle Part, North Arm, East Arm, Southeast Arm and Neck [4].

Gorontalo which is part of the North Arm of Sulawesi has very complex geological conditions due to tectonic influences. The spread of quarter age limestone in the Gorontalo plains and Pliocene-Pleistocene limestone around Lake Limbo to are evident. The effect of tectonics is very strong, especially from the position of reef limestone that is quarter age in the Gorontalo region. This reef limestone undergoes a very strong elevation, proven by field data near Gorontalo and the northern coast of Tanjung Daka [4-6].

The influence of tectonics on uplift rate limestone in Gorontalo has yet to be known how fast until now. This is inseparable from the data of limestone of Gorontalo that is regionally by dividing it into two large formations without knowing the absolute age and paleobathimetry so that the uplift rate of Gorontalo limestone is unknown. For this reason, based on the background, three main objectives of this research were raised, which are, to find out the absolute age of limestones, to know paleobathimetry and uplift rate of limestone.

Material and Method. The research location is in Yosonegoro region, Gorontalo Regency, which is the northern part of the Lake Limbo to basin with coordinates (00°39'6.7222"North, 122°54'50.0385"East) to (00°39'6.9397" North, 122°54'57.5275"East) (figure 1). 24 meter high limestone outcrop research material. The research method carried out consisting of three methods namely the measured section (MS), biostratigraphy analysis and tectonic analysis. The MS method measures the detail of the coating using a Jacob's staff at 1.5 meter intervals by systematic lithology sampling from the oldest to the youngest rocks [7, 8]. Biostratigraphy analysis using the Olympus SZ61 binocular microscope to identify the types of fossil planktonic and benthonic for a mini ferain determining the age of rocks and paleobathimetry [9-19]. Tectonic analysis to calculate the uplift rate based on absolute age data of limestone, paleobathimetry and current position of elevation [20-22].




Figure 1 – Location of limestone research in Limboto area, Gorontalo Regency, Indonesia

Results and discussion. The measured section results at the research location helped to determine the thickness of one stratigraphy path way and determine the exact position of the sample for biostratigraphy analysis. A total of 10 samples that carried out biostratigraphy analysis based on facies obtained from measured section results. The samples taken certainly contain planktonic foraminifera fossils and benthic based on grain size and the content of foraminifera fossils.

Biostratigraphy analysis in 10 limestone samples contained 17 types of planktonic foraminifera fossil. The distribution of 17 types of planktonic foraminifera fossil can be seen in figure 2. Based on figure 2, interpretation and analysis of the distribution of the biostratigraphy zone and the position of the bio datum in the stratigraphy path way can be done. The research location is divided into two biostratigraphy zones, with details as follows:

1. Zone PL2

Definition: The upper boundary of PL2 zone is not found. The lower boundary is LO *Globoquadrina dehiscens*.

Discussion: This zone is *Globoquadrina dehiscens* partial range zone. Zone PL2 (part) is equivalent to the zone PL2 [14] and zone N19 [23]. This zone represents the youngest planktonic foraminifera (top). LO *Globoquadrina dehiscens* [14, 24] were found in 3G samples. The contents of fossil association in this zone are *Globigerinoides desimmaratus* (deLeroy), *Globigerinoides desquadrilobatus* (d'Orbigny), *Globigerinoides desruber* (d'Orbigny), *Globigerinoides desruber* (d'Orbigny), *Globigerinoides dessacculiferus* (Brady), *Globorotalia humerosa* (Takayanagi and Saito), *Globorotalia plesiotumida* (Blow and Banner), *Pulleniatina precursor* (Banner and Blow) and *Pulleniatina obliquiloculata* (Parker and Jones). In this sample also found rework fossils in the form of *Globorotalia juanai* (Bermudez and Bolli) and *Globorotalia menardii* (d'Orbigny in Parker, Jones and Brady). The presence of this species is estimated to be reworks originating from older rocks. The thickness of this zone is 13 meters (interval 11-24 meters)

Age: Early Pliocene. ? – 5,80Ma.

2. Zone PL1

Definition: The upper boundary is LO *Globoquadrina dehiscens*. The lower boundary is not found.

Discussion: This zone is *Globoquadrina dehiscens* partial range zone. Zone PL1 is equivalent to the zone PL1 [14] and zone N18 [23]. The initial datum of this zone is not found. At the end of this zone is LO *Globoquadrina dehiscens* [14, 24]. LO *Globoquadrina dehiscens* was found in 3G samples. The presence of *Globoquadrina dehiscens* in samples of 3B, 3C, 3D, 3E, 3F and 3G. This species was not found in the samples above, namely 3H, 3I and 3J samples.

The contents of fossil association in this zone are *Globigerinoides desimmaratus* (de Leroy), *Globigerinoides desquadrilobatus* (d'Orbigny), *Globigerinoides desquadrilobatus* (deOrbigny), *Globigerinoides desruber*

(d'Orbigny), *Globigerinoi dessaculiferus* (Brady), *Globigerinoi dessubquadratus* (Bronnimann), *Globorotalia acostaensis* (Blow), *Globorotalia humerosa* (Takayanagi and Saito), *Globorotalia tumidatumida* (Brady), *Hastigerinaa equilateralis* (Brady), and *Pullenia tinapracursor* (Banner and Blow). In this sample also found rework fossil in the form of *Globorotalia continuosa* (Blow) and *Globorotalia juanai* (Bermudez and Bolli). The presence of this species is estimated to be reworks originating from older rocks. The thickness of this zone is 11 meters (interval 1 - 11 meters).

Age: Early Pliocene. 5,80 - ? Ma.




Figure 2 – Distribution chart danbiozo nation planktonic foraminifera in research area

Based on the division of two biostratigraphy zones, the position of the bio datum of LO *Globoquadrina dehiscens* was found at a thickness of 11 meters, namely the 3G sample. Based on the classification of [14] the absolute age of LO *Globoquadrina dehiscens* was 5.80 Ma (million years ago). After knowing the absolute age, the interpretation of the position of the elevation location of the study and paleobathymetry were interpreted in the 3G sample.

The elevation position of the research location is based on direct measurements using the Global Positioning System(GPS) which is 70 meters.

The Paleobathymetry of 3G sample based on biostratigraphy analysis refers to two methods, namely the method of [9] and the combined method of [11, 25]. The Paleobathymetry of 3G sample refers to the method of [9] using the calculation of the formula equation $D = e^{(0.0061.P + 1.25)}$. D value is depth while P is the pelagic ratio. The requirement for using this formula equation is the P value must be known from the comparison of the content of the number of fossils of planktonic and benthic foraminifera. The P value in sample 3G after being calculated is 75% so that the value of D or paleobathymetry can be known as 338.661 meters.

The Paleobathymetry of 3G sample refers to the method of [11] based on the overlay of benthic foraminifera fossil content. The fossil content of benthic foraminifera consists of *Fijionionfijiense* (Cushman and Edwards), *Gyrodinoidessoldanii* (d'Orbigny), *Melonisaffinis* (Reuss), *Nonionfabum* (FichtelandMoll), *Pararotaliavenusta* (Brady), *Praeglobobulimina ovata* (d'Orbigny), *Rhabdammina-discreata* (Brady) and *Saccorhizaramosa* (Brady). Based on the content of benthic foraminifera fossil, paleobathymetry is obtained in the upper lobe zone with a depth of 283.65-366 meters. The Paleobathymetry obtained is still in the range of depth, then the paleobathymetry correction curve [25] was used by looking for the middle value (median) obtained 324.825 meters

Calculation of the uplift rate in the research area was carried out by comparing the total depth of the limestone since it was formed or deposited until this time divided by the absolute age of the limestone. The total depth value of the limestone summed the thickness of the bio datum position from MS result with the current elevation position and paleobathymetry. Calculation of limestone elevation using two data paleobathymetry namely the method of [9] and the combined method of [11,25]. The results of calculating the limestone uplift can be seen in table.

Calculation Comparison of Limestone Uplift Rate

Method	Paleobathymetry (meters)	Elevation (meters)	Bio datum position (meters)	Uplift rate (mm/year)	Rate comparison
[9]	338.661	70	11	0.0724	1 : 1
[11, 25]	324.825	70	11	0.0699	

Comparative analysis of the uplift rate of limestone using both methods produces almost the same results. Difference in the ratio of the two uplift rates if rounded 1: 1 or the uplift range 0.0699-0.0724 mm/year.

Conclusion. Biozonation of planktonic foraminifera in the research area consisted of two, namely PL1Zone and PL 2Zone with bio datum of LO *Globoquadrina dehiscens*(5.80Ma) equivalent to the Early Pliocene. The depth of limestone was calculated based on the calculation of the total position of the bio datum of MS results with the current elevation position and paleobathymetry. Based on the data of absolute age and limestone depth since it was formed until it was elevated up to its current position, the uplift rate of limestone in the study are arranged from 0.0699-0.0724 mm/year.

Acknowledge. I would like to thank *Educational Fund Management Agency* (LPDP) for helping me provide the educational scholarships through *Indonesian lecturer main scholarship* (BUDI DN).

Аанг Панджи Пермана^{1,2}, Сабаджио Прэмамиджой², Акмалуддин²

¹Горонтало мемлекеттік университеті, Горонтало, Индонезия,

²Гаджа Мада университеті, Джокьякарта, Индонезия

БИОСТРАТИГРАФИЯСЫН ТАЛДАУ НЕГІЗІНДЕ
ГОРОНТАЛО (ИНДОНЕЗИЯ) ӘҚТАСЫНЫҢ КӨТЕРІЛУ ЖЫЛДАМДЫҒЫ

Аанг Панджи Пермана^{1,2}, Сабаджио Прэмамиджойо², Акмалуддин²

¹Государственный университет Горонтало, Инженерно-геологический факультет,
Горонтало, Индонезия,

²Университет Гаджа Мада, Инженерно-геологический факультет,
Джокьякарта, Индонезия

СКОРОСТЬ ПОДЪЕМА ИЗВЕСТНЯКА ГОРОНТАЛО (ИНДОНЕЗИЯ) НА ОСНОВЕ АНАЛИЗА БИОСТРАТИГРАФИИ

Аннотация. Скорость подъема известняка Горонтало очень сильно зависит от полноты и глубины данных. Все необходимые данные могут быть получены на основе полного анализа биостратиграфии. В качестве исследовательского материала использовался слой известняка толщиной 24 м. Целью исследования явилось определение абсолютного возраста известняка, палеобатиметрии и скорости поднятия известняка на исследуемой территории. Используются три метода, состоящие из измерительного разреза (ИР), биостратиграфического анализа и тектонического анализа. В качестве исходного био-уровня в известняке использовался только *LO Globoquadrina dehiscens*. Определение палеобатиметрии проводилось с использованием двух методов. Скорость подъема известняка составила 0,0699-0,0724 мм/год.

Ключевые слова: скорость подъема, известняк, Горонтало, биостратиграфия.

Information about authors:

Aang Panji Permana, Gorontalo State University, Geological Engineering Department, Gorontalo, Indonesia; Universitas Gadjah Mada, Geological Engineering Department, Yogyakarta, Indonesia; aang@ung.ac.id; <https://orcid.org/0000-0002-6865-3564>

Pramumijoyo Subagyo, Universitas Gadjah Mada, Geological Engineering Department, Yogyakarta, Indonesia; bagyo@ugm.ac.id; <https://orcid.org/0000-0001-7524-1122>

Akmaluddin, Universitas Gadjah Mada, Geological Engineering Department, Yogyakarta, Indonesia; akmaluddin@gmail.com

REFERENCES

- [1] Hamilton W. (1979) Tectonics of the Indonesian region. Geological Survey Professional Paper 1078, U.S. Govern. Printing Office, Washington. U.S.G.S. Professional Paper 1078. P. 345.
- [2] Silver E.A., McCaffrey R., Smith R.B. (1983a) Collision, rotation and the initiation of subduction in the evolution of Sulawesi, Indonesia // J. Geophysic. Res. 88, B11: 9407-9418.
- [3] Hutchison C.S. (1989) Geological evolution of Southeast Asia. Oxford Monograph on Geology and Geophysics no 13, Oxford. P. 368.
- [4] Sukanto R., Ratman N. (2013) Quaternary rock, In :Surono., Hartono U (Eds), Sulawesi geology, Center for Surveying Geological Resources. LIPI Press: 110-112.
- [5] Katili J.A. (1970) Large transcurrent faults in southeast asia with special reference to Indonesia // International Journal of earth Science. 59 (2) : 581-600.
- [6] Bachri S., Partoyo E., Bawono S.S., Sukarna D., Surono., Supandjono J.B. (1997) Regional geology of Gorontalo, North Sulawesi. Collection of research and mapping results papers // Center for geological research and development: 18-30.
- [7] Compton R.R. (1985) Geology in the field. Wiley Press-New York. P. 416.
- [8] Bukhsianidze M., Chagelishvili R., Lordkipanidze D. (2018) Late Miocene Vertebrate Site of Chachunalori valley, Georgia, Southern Caucasus // Bulletin of the Georgian National Academy of Sciences. 12(2) : 70-75.
- [9] Van Marle L.J., Van Hinte J.E., Nederbragt A.J. (1987) Plankton percentage of the foraminiferal fauna in seafloor samples from the Australian-Irian Jaya continental margin, eastern Indonesia. Marine Geology. 77 : 151-156.
- [10] Van der Zwaan G.J., Jorissen F.J., De Stigter H.C. (1990) The depth dependency of planktonic/benthic foraminiferal ratios: constraints and applications. Marine Geology 95 : 1-16.
- [11] Jones R.W. (1994) The Challenger Foraminifera. Oxford: Oxford University Press. P. 149.
- [12] Chaisson W.P., Pearson P.N. (1997) Planktonic foraminifer biostratigraphy at site 925: middle miocene-pleistocene // In: Shackleton N.J., Curry W.B., Richter C., Bralower T.J. (Eds.). Proceedings of the Ocean Drilling Program, Scientific Results 154 : 3-31.
- [13] Nichols G. (2009) Sedimentology and Stratigraphy. London: Blackwell Science Ltd. P. 335.

- [14] Wade B.S., Pearson P.N., Berggren W., Palike H. (2011) Review and revision of cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale // *Earth Science Review*. 104 : 111-142.
- [15] Chunlian L., Yi H., Jie W., Guoquan Q., Tinting Y., Lianze X., Suqing Z. (2012) Miocene-pliocene planktonic foraminiferal biostratigraphy of the Pearl River Mouth Basin, Northern South China Sea // *Journal of Palaeogeography*. 1(1) : 43-56.
- [16] Ghosh A.K., Sarkar S. (2013) Facies analysis and paleoenvironmental interpretation of Piacenzian carbonate deposits from the Guitar Formation of Car Nicobar Island, India // *Geoscience Frontiers*. 4 : 755-764.
- [17] Martins M.V.A., Silva F., Laut L.L.M., Frontalini F., Clemente I.M.M.M., Miranda P., Figueira R., Sousa S.H.M., Dias J.A.M. (2015) Response of benthic foraminifera to organic matter quantity and quality and bioavailable concentrations of metals in Aveiro Lagoon (Portugal) // *PLoS ONE* 10 (2): 1-23. Doi:10.1371/journal.pone.0118077.
- [18] Roozpeykar A., Moghaddam I.M. (2016) Benthic Foraminifera as Biostratigraphical and Paleoecological Indicators: An Example from Oligo-Miocene Deposits in the SW of Zagros Basin, Iran // *Geoscience Frontiers* 7 : 125-140.
- [19] Oladimeji A., Adeyinka S.A., Adekeye O.A., Olesegun O., Emmanuel O.F. (2017) Foraminifera Biostratigraphy and Depositional Environment of Sediment in Sile Well Offshore Dahomey Basin Benin Republic // *Mayfeb Journal of Environmental Science*. 1 : 18-33.
- [20] Berggren W.A. (1992) Paleogene planktonic foraminifera magnetobiostratigraphy of the Southern Kerguelen Plateau (Sites 747-749) // In Wise S.W., Jr., Schlich R., et al. // *Proc. ODP, Sci. Results*, 120 (Pt. 2): College Station, TX (Ocean Drilling Program) : 551-568.
- [21] Barianto D.H., Kuncoro P., Watanabe K. (2010) The use of foraminifera fossils for reconstructing the Yogyakarta graben, Yogyakarta, Indonesia // *J. SE Asian Apl. Geol.* 2(2) : 138-143.
- [22] Shen C.C., Wu C.C., Dai C.F., Gong S.Y. (2018) Variable uplift rate through time: Holocene coral reef and neotectonics of Luta, eastern Taiwan // *Journal of Asian Earth Sciences*. 156 : 201-206.
- [23] Blow W.H. (1969) Late middle eocene to recent planktonic foraminiferal biostratigraphy // In: Brönnimann P., Renz H.H. (Eds.). *Proceedings of the First International Conference on Planktonic Microfossils*, E.J. Brill, Leiden 1 : 199-422.
- [24] Berggren W.A., Hilgen F.J., Langereis C.G., Kent D.V., Obradovich J.D., Raffi I., Raymo M.E., Shackleton N.J. (1995a) Late neogene chronology: new perspectives in high-resolution stratigraphy // *Geol. Soc. Am. Bull.* 107 : 1272-1287.
- [25] Van Hinte J.E. (1978) *Geohistory Analysis – Application of Micropaleontology in Exploration Geology* // *The American Association of Petroleum Geologists Bulletin*. 62 (2) : 201-222.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://www.geolog-technical.kz/index.php/en/>

Верстка Д. Н. Калкабековой

Подписано в печать 15.11.2019.
Формат 70x881/8. Бумага офсетная. Печать – ризограф.
19,7 п.л. Тираж 300. Заказ 6.