

**ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ФЫЛЫМ АКАДЕМИЯСЫНЫҢ
Satbayev University

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Satbayev University

N E W S

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Satbayev University

**SERIES
OF GEOLOGY AND TECHNICAL SCIENCES**

5 (443)

SEPTEMBER – OCTOBER 2020

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Бас редакторы
э. ф. д., профессор, КР ҮФА академигі
И.К. Бейсембетов

Бас редакторының орынбасары
Жолтаев Г.Ж. проф., геол.-мин. ф. докторы

Редакция алқасы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., академик (Қазақстан)
Абсадыков Б.Н. проф., корр.-мүшесі (Қазақстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қыргызстан)
Буктуков Н.С. проф., академик (Қазақстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Жарменов А.А. проф., академик (Қазақстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., академик (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«КР ҮФА Хабарлары. Геология және техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» РКБ (Алматы қ.).

Қазақстан Республикасының Акпарат және қоғамдық даму министрлігінің Акпарат комитетінде 29.07.2020 ж. берілген № KZ39VPY00025420 мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *геология және техникалық ғылымдар бойынша мақалалар жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2020

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыр көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
д. э. н., профессор, академик НАН РК
И. К. Бейсембетов

Заместитель главного редактора
Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н на я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., академик (Казахстан)
Абсадыков Б.Н. проф., чл.-корр. (Казахстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Буктуков Н.С. проф., академик (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Жарменов А.А. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., академик (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № KZ39VPY00025420, выданное 29.07.2020 г.

Тематическая направленность: *публикация статей по геологии и технических наукам.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2020

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief
doctor of Economics, professor, academician of NAS RK
I. K. Beisembetov

Deputy editor in chief
Zholtayev G.Zh. prof., dr. geol-min. sc.

Editorial board:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., academician (Kazakhstan)
Absadykov B.N. prof., corr. member. (Kazakhstan)
Agabekov V.Ye. academician (Belarus)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Buktukov N.S. prof., academician (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadzhikistan)
Gravis R.M. prof. (USA)
Zharmenov A.A. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., academician (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Ozdoyev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).
The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan **No. KZ39VPY00025420**, issued 29.07.2020.

Thematic scope: *publication of papers on geology and technical sciences.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 5, Number 443 (2020), 141 – 149

<https://doi.org/10.32014/2020.2518-170X.114>

UDC 551.4:571.6

**Zh. S. Mustafayev¹, A. T. Kozykeyeva¹, L. N. Ryskulbekova¹,
A. E. Aldiyarova¹, Arvydas Povilaitis²**

¹Kazakh National Agrarian University, Almaty, Kazakhstan;

²Vytautas Magnus University, Kaunas, Lithuania.

E-mail: z-mustafa@rambler.ru; aliya.kt@yandex.ru, ryskulbekova.laura@mail.ru,
ainur_005@mail.ru, arvydas.povilaitis@vdu.lt

GEOMORPHOLOGICAL ANALYSIS OF THE ILI RIVER BASIN CATCHMENT AREA FOR INTEGRATED DEVELOPMENT

Abstract. Based on long-term informational and analytical materials of the World Meteorological Organization (WMO), the Weather and Climate reference portal and stationary meteorological stations of the RSE Kazhydromet located in the catchments of the Ili River basin, which cover the Almaty region (Narynkol, Tekes, Sumbe, Dobyn, Aydarly, Kapshagay, Usharal, Bakanas, Kokzhide, Kuigan) of the Republic of Kazakhstan, Xinjiang Uygur Autonomous Region (Tekes, Xinyuan, Tokkuztara, Yamata, Kuldzh) of the People's Republic of China and using the law of geographic vertical zoning, the energy resources of river basins and groundwater, the climatic potential of natural systems that characterize the heat and moisture supply of natural landscapes and ecological and hydrogeochemical indicators showing the direction and intensity of the hydrogeochemical process on a spatial scale, which allowed for geomorphological zoning, are determined catchments of the Ili river basin, characterizing the natural functions of the river basin, to have runoff and environment formation, which are the basis for environmental management and environmental engineering.

Key words: climate, nature, resources, river, pool, environment, landscape, catchment, system, heat, moisture, security, potential, process, nature management, environmental management, function.

Relevance. The general current situation of the catchment of the basin of the transboundary Ili River, located on the territory of the People's Republic of China and the Republic of Kazakhstan, is characterized by a rather intense geoecological state. This situation is caused by the progressive involvement and development of the natural resource potential of vast territories (in our case, watersheds), the increase in technogenic impact on them and the disruption of the relationship between natural components in geosystems and in the human-nature system.

Therefore, to solve the geoecological problems of the watersheds of the transboundary river basin, it is necessary to consider and study them in the form of geosystems of a certain rank, including an interdependent set of components and developing as a whole on the basis of geomorphological schematization, describing the main processes of the functioning of watersheds with a possibly large set of parameters that take into account changes in the components of geosystems catchment areas.

The main natural function of the river basin is, firstly, drainage, secondly, peculiar combined geosystems (the principle of unification here is the unity of hydrogeochemical flows having one object for their discharge) and, thirdly, this is a spatial basis for nature management (allocation of land for various purposes) and environmental management, that is, objects of a comprehensive arrangement of their watersheds.

With a comprehensive arrangement of the catchments of the river basin, the catenary approach is the basis of the geomorphological schematization of the catena, which consists of four facies with different altitudinal positions. The eluvial facies represents a hill near the dividing line, the transeluvial facies represent the slope to the inflection point, the transaccumulative facies represent the slope after the

inflection point, and the supersquale is the lowland of the floodplain terraces. The floodplains of the rivers, despite their significance, were not considered in the work. The transeluvial and transaccumulative facies form the transit facies of the slope, and the supequal facies adjoins the watercourse.

Purpose of the study – based on geomorphological schematization of catchments of river basins using the catenary approach, zoning of the landscape system of catchments of the Ili River basin to identify regional differences and the need to accurately determine their economic and economic capabilities.

Object of study - The Ili River, which originates in the Central Tien Shan on the territory of the People's Republic of China (PRC) after the confluence of three tributaries - the rivers Kash, Kunes and Tekes, with the latter having a large tributary - the Koksu River and its flow formation zone (Halyktau and Narat) is characterized by the largest layer of precipitation - more than 1300 mm / year. The sources of the Tekes river are located on the territory of Kazakhstan on the Muzart glaciers and their length is 438 km, the catchment area is 28100 km² (within the Republic of Kazakhstan, respectively, 218 km and 4250 km²). The river, high in the upper reaches, at the intersection of the Tekes Depression loses a significant part of its flow for water consumption in the branches of the economy, but after the confluence of the tributaries Kokpak, Bayankol, Narynkol, which feed on meltwater from the Khan-Tengri glaciers, the water content of the river increases significantly. Then it flows through the territory of the People's Republic of China (East Turkestan), where it merges with the Kunes and Kash rivers, and at the 250-th km from their confluence enters the territory of Kazakhstan with the Ili river with many waters. After exiting the Kapshagai Gorge, the Ili River flows through the deserted Balkash plain to Lake Balkhash. Downstream the riverbed is unstable, replete with elders and islets. When it flows into the lake, the river forms a delta with an area of about 9000 km², which is divided into three systems of branches - Topar, Ili and Zhideli [1].

Research Methods and Materials. The work used information and analytical materials of the World Meteorological Organization (WMO), the reference and information portal «Weather and Climate» and stationary meteorological stations of the RSE «Kazhydromet» located in the catchments of the Ili River basin, which cover the Almaty region (Narynkol, Tekes, Sumbe, Dobyn, Aydarly, Kapshagay, Usharal, Bakanas, Kokzhide, Kuigan) of the Republic of Kazakhstan, Xinjiang Uygur Autonomous Region (Tekes, Xinyuan, Tokkuztara, Yamata, Kulja) of the People's Republic of China [2]

By the methodology of the comprehensive arrangement of the catchments of river basins, given the multidimensional nature of the problem, the whole set of methodological approaches that exist in environmental management, that is, the geosystem and catenary approaches, has been adopted.

The catenary approach is the basis of the geomorphological schematization of catenas when substantiating the need for a comprehensive arrangement of catchments of river basins [3,4,5].

In the geomorphological schematization of landscape catenas of catchment areas, in order to justify the need to equip river basins, each catchment within the same physical and geographical area is represented by a catena consisting of four facies with different altitude positions, determined by the depth of the relief: eluvial, transit, superaqual and subaqueous. If the watersheds have extended slopes, the transit facies is divided into the transeluvial and transaccumulative facies [5].

Thus, the basis of geomorphological schematization of watershed landscapes based on the geosystem approach, based on the excess of the earth's surface over the watercourse, makes it possible to more objectively differentiate the zones of facies locations.

However, in the quantitative description of the hydrogeochemical processes of catchments of river basins, the main integral factor is the energy of the groundwater flow, that is, the speed of their movement, the mass of water, depending on the thickness of the aquifer, filtration energy, and evaporation energy, which are not taken into account when using the excess of the earth's surface over the watercourse as a criterion for geomorphological schematization of river basins, i.e Δ_i and ΔH characterize the excess of the earth's surface over the watercourse, then you can imagine the energy or work performed by the flow of groundwater in the following form [4]: $\Delta E = A_i = m_i \cdot g \cdot \Delta H = m_i \cdot g \cdot \Delta_i$, where ΔE – change in energy at the site dx , kJ; A_i – work performed in an elementary volume by the flow of groundwater in the area dx , kJ; m_i – groundwater mass; Δ_i и ΔH – height exceeding the Earth's surface above the bank of the watercourse, m.

Construct geomorphological circuit landscape catenas possible using hydrogeochemical potential groundwater flow river basin (\bar{M}), characterizing work (\bar{A}_n), committed by a liquid in the process of precipitation to the ratio of the concentration of soil solution (\bar{C}_n), that is, they can be considered as the ability of the soil cover to be released from readily soluble salts from the upper to the lower reaches of the river basins: $\bar{M} = \bar{A}_n / \bar{C}_n$, where: \bar{M} – hydrogeochemical potential of river basins; \bar{A}_n – work performed in an elementary volume by the flow of infiltration water in the soil layer of river basins; \bar{C}^* – average salt concentration in groundwater flow [4]: ε

$$\bar{A}_n = O_c / \left[\frac{R}{L} - (1-t) \frac{R}{L} (1-\bar{\Delta}) \right], \quad \bar{C}^* = \left[C_o + (1-t) \frac{R}{L} (1-\bar{\Delta}) \cdot C_r / O_c \right] / C_{don},$$

where: R – radiation balance; O_c – precipitation; L – latent heat of vaporization; C_o – initial concentration of soil solution in the soil layer; $-$ permissible concentration of salts in the soil solution, which corresponds to the parameter of non-saline soils; C_r – salt concentration in groundwater; $(1-t)$ – infiltration action time ($t = T/365$), T – duration of the biological active period; $\bar{\Delta} = \Delta / \Delta_{kp}$, Δ – groundwater depth; Δ_{kp} – critical groundwater depth.

Research results. For the geomorphological zoning of the watershed territories of the Ili River basin, a catenary approach was used, which involves a geomorphological schematization of landscape catenas of the watersheds of river basins characterizing landscapes in the mountain class (eluvial facies), piedmont landscape subclass (transelyuvial facies), foothill plain landscape subclass (transaccus) flat class of landscapes (super-aquatic and sub-aquatic facies).

Based on the above methodological approach, the energies and work performed by the groundwater flow in the catchments of the Ili River basin are determined (table 1).

Table 1 – Energy resources of the catchments of the Ili River basin

Natural and climatic zones		Weather stations	The absolute height of the earth's surface, m	ΔH , m	Groundwater flow energy, kJ	
landscape class	facies				A_i	ΣA_i
Mount	Eluvial	Basuarkor	4059,0	–	–	–
		Narynkol	1806,0	1797,0	17628,6	31961,3
		Tekes	1766,0	40,0	392,4	14333,3
Piedmont	Trans-eluvial	Sumbe	1232,0	534,0	5238,5	13940,3
		Tekes (China)	1203,0	29,0	284,5	8701,8
		Xinyuan (China)	947,0	256,0	2511,4	8417,3
		Tokkuztara (China)	773,0	174,0	1707,0	5905,9
Piedmont lowland	Trans accumulative	Yamato (China)	723,0	50,0	490,5	4198,9
		Gulja (China)	663,0	60,0	588,6	3708,4
		Kokdala (China)	627,0	36,0	353,2	3119,8
		Dubyn	596,0	31,0	304,1	2766,6
		Aydarly	576,0	20,0	196,2	2462,5
		Kapshagay	540,0	36,0	353,2	2266,3
Plain	Super aqua	Ili	490,0	50,0	490,5	1913,1
		Bakbakty	459,0	31,0	304,1	1422,6
		Usharal	397,0	62,0	608,2	1118,5
		Bakanas	396,0	1,0	9,8	510,3
		Akdala	390,0	6,0	58,9	500,5
		Akkol	384,0	6,0	58,9	441,6
	Subaquatic	Araltobe	357,0	27,0	264,9	382,7
		Kokzhide	350,0	7,0	68,7	117,8
		Kuigan	345,0	5,0	49,1	49,1

As can be seen from table 2, in the catchment areas of the Ili River basin in the territory of the Raiymbek district of Almaty region, that is, the zone of runoff formation, where the mountain class of landscapes with eluvial facies is located, has rather high energy resources of river basins and groundwater, which are 23259.5 kJ. At the same time, in the watershed territories of the Ili River basin, which pass through the borders of the Xinjiang Uygur Autonomous Region of the People's Republic of China, are located in the piedmont class of landscapes with transeluvial facies, the energy resources of river basins and groundwater are reduced to 8034.4 kJ. The energy resources of river basins and groundwater in the foothill lowland class of landscapes with a transaccumulative facies of catchments of the Ili River basin, which cover part of the territory of the Xinjiang Uygur Autonomous Region of the People's Republic of China and the Uyghur District of the Almaty Region of the Republic of Kazakhstan, amount to 2285.8 kJ. At the same time, in the territories of the catchment part of the Republic of Kazakhstan, covering from the Dobyn hydrological post to the city of Kapshagai, it amounts to 500.3 kJ, that is, the energy resources of the river basin and groundwater are sharply reduced. In the territories of the flat class of landscapes with superaqual facies, covering the Ili and Balkhash districts of the Almaty region, the energy resources of the river basin and groundwater are reduced to 1530.3.1 kJ, and in the zone of the subaqueous facies it decreases sharply to 333.6 kJ.

Thus, in accordance with the principle of geographical zonality existing in energy and groundwaters, from 31,961.3 to 49.1 thousand catchment areas of the Ili River basin (table 2).

Table 2 – Geomorphological schematization of landscape catenas in the catchment areas of the Ili River basin according to the energy resources of the river basin and groundwater

Natural and climatic zones		Geomorphological indicator	
Landscape class	Facies	The absolute height of the earth's surface, m	Groundwater flow energy, kJ
Mount	Eluvial	<1800	<175000
Piedmont	Trans-eluvial	800-1800	8000-17500
Piedmont lowland	Trans accumulative	540-800	5500-8000
Plain	Super aqua	350-540	3500-5500
	Subaqueous	>350	>3500

The natural heat and moisture supply of the active surface is the most important constituent element of the complex of natural productive forces, that is, natural landscapes that are actively involved in the biological process in general and in the formation of natural-technogenic complexes, especially the catchments of river basins [6,7,8].

Based on the long-term information and analytical materials of the located meteorological stations in the catchments of the Ili River basin, their average long-term energy resources and natural and climatic potentials are determined, that is (table 2): the sum of the air temperature for the biologically active period of the year ($\Sigma t, {}^{\circ}\text{C}$), amount of precipitation ($O_c, \text{ mm}$), evaporation level ($E_o, \text{ mm}$), and photosynthetically active radiation ($R, \text{ kJ/cm}^2$) (table 3).

As can be seen from table 3, the energy resources of the watersheds of the Ili River basin from the mountainous class of landscapes (eluvial facies) to the plain class of landscapes (subaqueous) increase, the sum of the biological active air temperature from 2338.0 to 3800.0 ° C, volatility from 844, 0 to 1472.0 mm and the radiation balance from 134.0 to 182.0 kJ/cm^2 , and precipitation decreases from 433.0 to 144.0 mm. Under these conditions, landscape-geochemical catenes are formed in the catchment areas of the river basin, that is, the simplest cascade landscape-geochemical system, characterized by the intensity of hydrogeochemical flows, which largely depend on their energy resources.

Table 3 – Natural and energy resources of the catchments of the Ili River basin

Natural and climatic zones (landscape class, facies)	Weather stations	Absolute height (H), m	Natural and climatic indicators			
			O_c , mm	$\sum t$, °C	E_o , mm	R , kJ/cm ²
Mountain class of landscapes (eluvial facies)	Narynkol	1806,0	433,0	2805,0	1001,0	149,0
	Tekes	1766,0	421,0	2338,0	844,0	134,0
Piedmont subclass of landscapes (transeluvial facies)	Sumbe	1232,0	377,0	3074,0	1122,0	158,0
	Tekes (China)	1203,0	259,0	3100,0	954,0	159,0
	Tokkuztara (China)	773,0	248,0	3579,0	894,0	175,0
Piedmont lowland subclass of landscapes (trans accumulative facies)	Yamato (China)	723,0	252,0	3130,0	1269,0	162,0
	Gulja (China)	663,0	248,0	3800,0	1284,0	182,0
	Zharkent	641,0	213,0	3950,0	1661,0	187,0
	Dobyn	596,0	226,0	4100,0	1748,0	192,0
	Aydarly	576,0	364,0	4305,0	1247,0	199,0
	Kapshagay	540,0	370,0	3750,0	1528,0	180,0
Plain class of landscapes (superaqual facies)	Usharal	397,0	354,0	3622,0	1168,0	176,0
	Bakanas	396,0	273,0	3700,0	1527,0	179,0
Plain class of landscapes (subaqueous)	Kokzhide	350,0	224,0	3700,0	1474,0	179,0
	Kuigan	345,0	144,0	3800,0	1472,0	182,0

At the same time, the energy resources of river basins and groundwaters of the natural system are additionally characterized by hydrogeochemical processes of groundwater, which are characterized by ecological and hydrogeochemical parameters of landscape of river basins, which are extremely important in the ecological and hydrogeochemical zoning of landscape and geographical zones and water management assessment of catchments of the Ili River basin (table 4).

Table 4 – Environmental-hydrogeochemical indicator
of the landscape system of the catchment area of the Ili River basin

Natural and climatic zones (landscape class, facies)	Weather stations	H , m	Indicators				
			C_o , g/l	C_z , g/l	Δ , m	\bar{A}_n	\bar{C}^*
Mountain class of landscapes (eluvial facies)	Basuarkor	4059,0	0,30	1,00	10,0	0,73	0,60
	Narynkol	1806,0	0,30	1,00	10,0	0,79	0,60
	Tekes	1766,0	0,30	1,00	10,0	0,60	0,60
Piedmont subclass of landscapes (transeluvial facies)	Sumbe	1232,0	0,40	1,20	10,0	0,60	0,80
	Tekes (China)	1203,0	0,40	1,30	10,0	0,41	0,80
	Tokkuztara (China)	773,0	0,40	1,30	10,0	0,35	0,80
Piedmont lowland subclass of landscapes (trans accumulative facies)	Yamato (China)	723,0	0,50	1,50	6,0	0,39	1,00
	Gulja (China)	663,0	0,50	1,50	6,0	0,29	1,00
	Zharkent	641,0	0,50	1,50	6,0	0,34	1,00
	Dobyn	596,0	0,50	1,50	6,0	0,29	1,00
	Aydarly	576,0	0,50	1,50	6,0	0,46	1,00
	Kapshagay	540,0	0,50	1,50	6,0	0,51	1,00
Plain class of landscapes (superaqual facies)	Usharal	397,0	0,90	3,50	3,0	0,50	1,50
	Bakanas	396,0	0,90	3,50	3,0	0,38	2,50
Plain class of landscapes (subaqueous)	Kokzhide	350,0	1,50	6,00	3,0	0,31	1,80
	Kuigan	345,0	1,50	6,00	3,0	0,20	1,80

In this case, the work, as can be seen from table 4, performed in an elementary volume by a flow of infiltration water in the soil layer (\bar{A}_n) from the mountain side (eluvial facies) to the lowland (subaqueous) zones gradually decreases from 0.73 to 0.20, and the average salt concentration in surface water - soil - "groundwater" system (C^*), on the contrary, increases from 0.60 to 1.20. Consequently, the ecological-hydrogeochemical potential or soil-reclamation indicator of the landscape (M), obeying the law of vertical zonality, decreases from 1.21 to 0.13, which characterizes a saline collection basin in the area of hydrogeochemical flow stacking. This regularity shows the available possibilities for the formation of hydrogeochemical processes in the catchment areas of the Ili River basin, which are damaged by the spread in the flat class of landscape with superaqueous and suaqueous facies, where salinization of soils with low ecological productivity are widespread [57], which are estimated using integrated criteria for environmental and climatic productivity natural landscapes (table 5):

- humidification coefficient ($K_y = O_c / E_o$), where O_c – precipitation, mm; E_o – volatility over the biological active period of the year, mm: $E_o = 0.0018 \cdot (25 + t)^2 (100 - a)$, where t – is the average monthly temperature °C; a – monthly average relative humidity, % [9];

- dryness index ($\bar{R} = R / LO_c$, where L – specific heat of vaporization, assumed constant and equal to 2,5 kJ/cm²); R – photosynthetically active radiation, kJ/cm²[10]: $R = 13.39 + 0.0079 \cdot \sum t > 10^\circ\text{C}$, here $\sum t, ^\circ\text{C}$ – the sum of the air temperature for the biological active period of the year [11].

At the same time, under natural conditions, the formation of landscape hydrogeochemical facies largely depends on the heat and moisture supply of the catchment areas of the Ili River basin, since the natural humidity coefficient decreases from 0.50 from the mountain class of landscapes (eluvial facies) to the Plain class of landscapes (subaqueous) to 0.10, and the "dryness index", which characterizes the degree of balance of moisture and heat, rises from 1,273 to 5,056, which show the presence of sufficiently high energy resources.

Table 5 – Geomorphological schematization of landscape catenas of catchments of the Ili River basin

Humidity zones	Indicators of heat and moisture			The absolute height of the earth's surface, m	Administrative districts
	Weather stations	K_y	\bar{R}		
Mountain class of landscapes (eluvial facies)					
Wet mountain	Narynkol	0,43	1,370	1806,0	Raiymbek
	Tekes	0,50	1,273	1766,0	
Piedmont subclass of landscapes (transeluvial facies)					
Arid mountain	Sumbe	0,34	1,676	1232,0	Raiymbek
	Tekes (China)	0,27	2,456	1203,0	Ili Kazakh Autonomous Prefecture
	Tokkuztara (China)	0,29	2,822	773,0	
Piedmont lowland subclass of landscapes (trans accumulative facies)					
Arid piedmont	Yamato (China)	0,20	2,571	723,0	Ili Kazakh Autonomous Prefecture
	Gulja (China)	0,19	2,935	663,0	
	Zharkent	0,13	3,510	641,0	Panfilov
	Dobyn	0,13	3,398	596,0	Uigur
	Aydarly	0,29	2,187	576,0	Kerbulak
	Kapshagay	0,24	1,946	540,0	Kapshagay city
Plain class of landscapes (superaqueous facies)					
Arid plain	Usharal	0,30	1,988	397,0	Ili
	Bakanas	0,18	2,622	396,0	Balkhash
Plain class of landscapes (subaqueous)					
Very arid desert	Kokzhide	0,15	3,200	350,0	Balkhash
	Kuigan	0,10	5,056	345,0	

Thus, using the laws of geographic vertical zonality, the energy resources of river basins and groundwater, the climatic potential of natural systems characterizing the heat and moisture supply of natural landscapes, and ecological and hydrogeochemical indicators showing the direction and intensity of the hydrogeochemical process on a spatial scale, which allowed geomorphological zoning of catchments of the Ili River basin, characterizing the nature valued functions of the river basin, ie runoff and environment formation and is a basis for environmental management and environmental engineering (table 5).

Thus, based on the geomorphological schematization of the catchments of the transboundary Ili river, based on the drainage of river basins as a thermodynamic system, the boundary of which coincides with the altitudinal zonality, it is possible to assess changes in the hydrological, ecological, soil, and ecological-hydrogeochemical regime of landscape systems taking into account natural and territorial differences as performing important environment formation or environmental functions.

Conclusions. Based on long-term informational and analytical materials of the World Meteorological Organization (WMO), the Weather and Climate reference portal and stationary meteorological stations of the RSE Kazhydromet, located in the catchments of the Ili River basin, which cover the Almaty region (Narynkol, Tekes, Sumbe, Dobyn, Aydarly, Kapshagay, Usharal, Bakanas, Kokzhide, Kuigan) of the Republic of Kazakhstan, Xinjiang Uygur Autonomous Region (Tekes, Xinyuan, Tokkuztara, Yamata, Kuldzh) of the People's Republic of China and using the law of geographic vertical zoning, the energy resources of river basins and groundwater, the climatic potential of natural systems that characterize the heat and moisture supply of natural landscapes and ecological and hydrogeochemical indicators showing the direction and intensity of the hydrogeochemical process on a spatial scale, which allowed for geomorphological zoning, are determined catchments of the Ili river basin, characterizing the natural functions of the river basin, to have runoff and environment formation, which are the basis for environmental management and environmental engineering.

**Ж. С. Мұстафаев¹, Ә. Т. Қозыкеева¹, Л. М. Рысқұлбекова¹,
А. Е. Алдиярова¹, Арвидас Повилайтис²**

¹Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан;

²Ұлы Витовт университеті, Каунас, Литва

КЕШЕНДІ ҮЙЛЕСТИРУ ҮШІН ІЛЕ ӨЗЕНІНІҢ СУ ЖИНАУ АЛАБЫН ГЕОМОРФОЛОГИЯЛЫҚ ТАЛДАУ

Аннотация. Қазақстан Республикасы Алматы облысының (Нарынкол, Текес, Сүмбе, Добын, Айдарлы, Қапшагай, Ушарал, Бақанас, Кокжиде, Құйған) және Қытай Халық Республикасының Синьцзян-Үйғыр автономиялық ауданының аймағын қамтитын Іле өзенінің су жинау алабына орналасқан «Қазгидромет» РМӘ-нің тұракты метеорологиялық бекеттерінің және Әлемдік метеорологиялық ұжымның (ӘМҰ) «Ауа-райы және климат» анықтамалық-ақпараттық жүйесінің көпжылдық мәліметі негізінде және географиялық тік белдеу заңын пайдалана отырып, өзен алабының және жерасты сұзы ағынының энергетикалық ресурстары, табиғи жүйенің табиғи-климаттық потенциалын сипаттайтын табиғи ландшафттардың ылғалмен қамтамасыз етілу көрсеткіштері, гидрохимиялық үдерістер бағытын және қарқының бейнелейтін уақыт-көңістік масштабындағы экологиялық-гидрохимиялық көрсеткіштері анықталды, ал ол оның нәтижесі өзен алабының табиғи қызметін сипаттайтын Іле өзенінің су жинау алабында геоморфологиялық ауданды жүргізуге мүмкіндік берді, яғни оның су ағынын және ортаны құруышы қасиеті, табигатты пайдалану және табигаты үйлестірудің негізі болып саналады.

Іле өзенінің су жинау алабы аймағының табиғи энергетикалық ресурстары ландшафттардың таулы топтарынан (элювиалдық фациядан) ландшафттардың жазықтық тобына (субаквальдық фацияға) қарай өседі, яғни биологиялық белсенді ауа температурасының жиынтығы 2338,0 °C-тан 3800,0 °C-қа, буланудың шамасы 844,0 мм-ден 1472,0 мм-ге және радиациялық тенгерме 134,0 кДж/см²-тан 182,0 кДж/см²-қа дейін, ал атмосфералық жауын-шашын шамасы 433,0 мм-ден 144,0 мм-ге дейін төмөндейді және осы жағдайға тән ландшафттық-геохимиялық катен, яғни геохимиялық ағын қарқыны арқылы ерекшеленетін, географиялық аймақтық (белдеулік) заңдылығына байланысты сатыланған ландшафттық-геохимиялық жүйе пайда болады және көптеген жерасты сұзы ағынының энергетикалық қуаты соган байланысты болғандықтан, оның сандық шамасы элювиалдық фациядан субаквальдық фацияға қарай 31961,3 кДж-дан 49,1 кДж-ға дейін біртіндеп төмөндейді, ал ол өзен аймағын геоморфологиялық жөлілеуге мүмкіндік береді.

Сонымен, табиғи ортандың нактыланған көлемдегі топырак қабатындағы сүзілген су ағынының атқаралын жұмысын (\bar{A}_n) таулы (элювиалдық фациядан) жазықтық (субаквальдик фацияға) аймаққа қарай 0,73-тен 0,20 -ға дейін төмендесе, «жер беті сұзы-топырак-жерасты сұзы» жүйесіндегі су ағынының орташа тұздылығы 0,60-тан 1,80-ға дейін өседі. Сондықтан, географиялық аймақтық (белдеулік) зандылығына байланысты экологиялық-гидрогохимиялық немесе топырак-мелиоративтік көрсеткіш (\bar{M}) 1,21-ден 0,11-ге дейін төмендейді, ал ол гидрогохимиялық ағынының қорлану аймағында тұз жиналатын алаптың қалыптасатынын сипаттайды, яғни бұл зандылық Іле өзенінде су жинау алабының аймағындағы гидрогохимиялық үдерістердің қалыптасу мүмкіншілігіне байланысты, оның жазықтық аймағында супераквальдик фациялардың таралыуына дәлел бола алады.

Сонымен катар, табиғи жағдайдағы ландшафттық-гидрогохимиялық фацияларының қалыптасуына Іле өзенінде су жинау алабының жылу және ылғалмен қамтамасыз ету дәрежесіне тікелей байланысты, себебі ландшафттардың таулы топтарынан (элювиалдық фациядан) ландшафттардың жазық топтарына (субаквальдик фацияға) қарай табиғи ылғалдану көрсеткіші 0,50-ден 0,10-ға дейін төмендей бастайды, ал жылу және ылғалдың тенгермелік дәрежесін сипаттайтын «кургақшылық белгісі» 1,273-ден 5,056-ға дейін жоғарылады, ал будай жағдай аймақтың энергетикалық ресурстарының өте жоғары екендігін көрсетеді.

Сонымен, өте маңызы ортанды құруышы және экологиялық қызмет атқаратын өзенінде су жинау алабы су ағынының қалыптасу жағдайын термодинамикалық жүйе ретінде қарастыра отырып, шекаралық Іле өзенінде су жинау алабын геоморфологиялық желілеу негізінде және аймақ шекарасының биік белдеулікке сай келуі, оның ландшафт жүйесінде гидрологиялық, экологиялық-топырак және экологиялық-гидрогохимиялық тәртібінің өзгеруін, табиғи-аймақ өзгешелігін ескере отырып бағалауға мүмкіндік береді.

Түйін сөздер: климат, табигат, қор, өзен, алап, орта, ландшафт, су жинау, жүйе, жылу, ылғал, қамтамасыз ету, потенциал, үдеріс, табигатты пайдалану, табигат үйлестіру, қызмет.

**Ж. С. Мустафаев¹, А. Т. Козыкеева¹, Л. М. Рысқулбекова¹,
А. Е. Алдиярова¹, Арвидас Повилайтис²**

¹Казахский национальный аграрный университет, Алматы, Казахстан;

²Университет Витовта Великого, Каунас, Литва

ГЕОМОРФОЛОГИЧЕСКИЙ АНАЛИЗ ВОДОСБОРА БАССЕЙНА РЕКИ ИЛИ ДЛЯ КОМПЛЕКСНОГО ОБУСТРОЙСТВА

Аннотация. На основе многолетних информационно-аналитических материалов Всемирной метеорологической организации (ВМО), справочно-информационного портала «Погода и климат» и стационарных метеорологических станций РГП «Казгидромет», расположенных в водосборах бассейна реки Или, которые охватывают Алматинскую область (Нарынкол, Текес, Сумбе, Добын, Айдарлы, Капшагай, Ушарал, Баканас, Кокжиде, Куйган) Республики Казахстан, Синьцзян-Уйгурский автономный район (Текес, Синьюань, Токкузтара, Ямату, Кульджа) Китайской Народной Республики и с использованием закона географической вертикальной зональности, определены энергетические ресурсы речных бассейнов и подземных вод, природно-климатического потенциала природных систем, характеризующих тепло- и влагообеспеченности естественных ландшафтов и эколого-гидрогохимических показателей, показывающих направленность и интенсивность гидрогохимического процесса в пространственном масштабе, которые позволили провести геоморфологическое районирование водосборов бассейна реки Или, характеризующих природные функции речного бассейна, то есть стокообразование и средообразование, являющихся базисом для природопользования и природобустроства.

Природные энергетические ресурсы территории водосборов бассейна реки Или от горного класса ландшафтов (элювиальная фация) в сторону равнинного класса ландшафтов (субаквальная) повышается, то есть сумма биологической активной температуры воздуха от 2338,0 до 3800,0 °C, испаряемость от 844,0 до 1472,0 мм и радиационный баланс от 134,0 до 182,0 кДж/см², а атмосферные осадки уменьшаются от 433,0 до 144,0 мм и в этих условиях в водосборных территориях речного бассейна формируется ландшафтно-геохимические катены, простейшая каскадная ландшафтно-геохимическая система, отличающаяся интенсивностью гидрогохимических потоков, которые во многом зависят от их энергетических ресурсов подземных вод, подчиняющихся от закона географической зональности, от элювиальной до субаквальной фации постепенно уменьшается от 31961,3 кДж до 49,1 кДж, что дает возможность на их основании производить геоморфологическую схематизацию ландшафтных катенов водосборной территории бассейна реки Или.

При этом работа, совершающаяся в элементарном объеме потоком инфильтрационных вод в почвенном слое (\bar{A}_n) от стороны горных (элювиальная фация) к равнинным (субаквальная) зонам, постепенно уменьшается от 0,73 до 0,20, а средняя концентрация солей в системе «поверхностная вода - почва - грунтовая вода» (\bar{C}^*), наоборот, увеличивается от 0,60 до 1,80. Следовательно, эколого-гидрогохимический потенциал

или почвенно-мелиоративный показатель ландшафта (\bar{M}), подчиняясь закону вертикальной зональности, уменьшается от 1,21 до 0,11, что характеризует в области магазинирования гидрогоеохимического потока формируется солесборный бассейн, то есть эта закономерность показывает имеющиеся возможности формирования гидрогоеохимических процессов в территориях водосбора бассейна реки Или, что повреждаются распространением в равнинном классе ландшафта с супераквальной и суаквальной фациями.

При этом в естественных условиях формирования ландшафтно-гидрогоеохимических фаций во многом зависят от тепло- и влагообеспеченности территорий водосборов бассейна реки Или, так как от горного класса ландшафтов (элювиальная фация) до равнинного класса ландшафтов (субаквальная) коэффициент естественного увлажнения уменьшается от 0,50 до 0,10, а «индекс сухости», характеризующий степень сбалансированности влаги и тепла, повышается от 1,273 до 5,056, которые показывают наличие достаточно высоких энергетических ресурсов.

Таким образом, на основе геоморфологической схематизации водосборов трансграничной реки Или, базирующейся на положениях стокообразования речных бассейнов как термодинамической системы, граница которых совпадает с высотной поясностью, позволяют оценить изменения гидрологического, экологопочвенного и эколого-гидрогоеохимического режима ландшафтных систем с учетом природно-территориальных различий как выполняющие важные средообразующие или экологические функции.

Ключевые слова: климат, природа, ресурсы, река, бассейн, среда, ландшафт, водосбор, система, тепло, влага, обеспеченность, потенциал, процесс, природопользование, природообустройство, функция.

Information about authors:

Mustafayev Zhumakhan Suleimenovich, Doctor of Technical Sciences, Professor, Professor of the Department «Water Resources and Melioration», Kazakh National Agrarian University, Almaty, Kazakhstan; z-mustafa@rambler.ru; <https://orcid.org/0000-0003-2425-8148>

Kozykeyeva Aliya Tobazhanovna, Doctor of Technical Sciences, Associate Professor, Professor of the Department «Water Resources and Melioration», Kazakh National Agrarian University, Almaty, Kazakhstan; aliya.kt@yandex.ru; <https://orcid.org/0000-0003-0581-0881>

Ryskulbekova Laura Meldakhanovna, Document PhD of the Department «Water Resources and Land Reclamation», Kazakh National Agrarian University, Almaty, Kazakhstan; ryskulbekova.laura@mail.ru; <https://orcid.org/0000-0002-1374-5920>

Aldiyarova Ainur Esirkepovna, PhD, Senior Lecturer of the Department of «Water Resources and Melioration», Kazakh National Agrarian University, Almaty, Kazakhstan; ainur_005@mail.ru; <https://orcid.org/0000-0002-6017-5182>

Arvydas Povilaitis, doctor of technical sciences, professor, Vytautas Magnus University, Kaunas, Lithuania; arvydas.povilaitis@vdu.lt; <https://orcid.org/0000-0003-1285-4604>

REFERENCES

- [1] Burlibaev M.Zh., Amirgaliev N.A., Shenberger I.V., Skolsky V.A., Burlibaeva D.M., Uvarov D.V., Smirnova D.A., Efimenko A.V., Milyukov D.Yu. Problems of pollution of the main transboundary rivers of Kazakhstan, Almaty: «Kanagat», 2014. Vol. 1. 744 p. (in Russ.).
- [2] Mustafayev Zh.S., Arvidas Povilaitis, Ryskulbekova L.N. Assessment of the natural and climatic potential of the catchment area of the Ili River // Studies, Results, 2019. N 1. P. 103-111 (in Russ.).
- [3] Kireicheva L.V., Kozykeeva A.T., Dauletbay S.D. Complex arrangement of the Shu River (monograph), Saarbrucken. Deutschland. 2016. 140 p. (in Russ.).
- [4] Mustafaev Zh.S. Kozykeeva AT, Zhidkova G.E., Dauletbay S.D., Zhanykhan K. Applied model of geomorphological schematization of landscape systems of river watersheds // International Technical and Economic Journal, 2016. N 3. P. 59-65 (in Russ.).
- [5] Golovanov A.I., Sukharev Yu.I., Shabanov V.V. Complex arrangement of territories - a further stage of land reclamation // Land reclamation and water management, 2006. N 2. P. 25-35 (in Russ.).
- [6] Adilbektegi G.A., Mustafayev J.S., Uvatayeva T.K., Dulatbekova Z.N., Jozef Mosiej. A new approach to the evaluation of bioclimatic potential of landscapes on the example of northern Kazakhstan // News of the National Academy of sciences of the Republic of Kazakhstan. Series of geology and technical sciences. ISSN 2224-5278. Vol. 5, N 437 (2019). P. 16-25 (in Eng.).
- [7] Adilbektegi G.A., Mustafayev J.S., Uvatayeva T.K., Dulatbekova Z.N., Jozef Mosiej. Quantitative and qualitative assessment of biological and ecological potential of the landscapes of southern Kazakhstan // News of the National Academy of sciences of the Republic of Kazakhstan. Series of geology and technical sciences. ISSN 2224-5278. Vol. 6, N 438 (2019). P. 96-103 (in Eng.).
- [8] Mustafayev Zh.S., Kozykeyeva A.T., Kalmashova A.N., Aldiyarova A.E., Arvydas Povilaitis. Ecological and water economic assessment of the Yesil river basin catchment area // News of the National Academy of sciences of the Republic of Kazakhstan. Series of geology and technical sciences. ISSN 2224-5278. Vol. 2, N 440 (2020), P. 123-131.
- [9] Ivanov N.N. Humidification zones of the globe // Izv. USSR Academy of Sciences. Series geography and geophysics. 1941. N 3. P. 15-32 (in Russ.).
- [10] Budyko M.I. Thermal balance of the earth's surface. L.: Gidrometeoizdat, 1956. 255 p. (in Russ.).
- [11] Nikolsky Yu.N., Shabanov V.V. Calculation of design productivity depending on the water regime of reclaimed land // Hydrotechnics and land reclamation. 1986. N 9. P. 52-56 (in Russ.).

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://www.geolog-technical.kz/index.php/en/>

Редакторы *M. С. Ахметова, Д. С. Аленов, А. Ахметова*
Верстка *Д. А. Абдрахимовой*

Подписано в печать 14.10.2020.
Формат 70x881/8. Бумага офсетная. Печать – ризограф.
14 п.л. Тираж 300. Заказ 5.